



Olfactory receptor 10C1 Monoclonal Antibody

Catalog No	BYmab-13438
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	OR10C1
Protein Name	Olfactory receptor 10C1
Immunogen	The antiserum was produced against synthesized peptide derived from human OR10C1. AA range:61-110
Specificity	Olfactory receptor 10C1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10C1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	34kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Testicle,
Function	function:Odorant receptor .,polymorphism:A stop codon at position Gln-55 in the gene coding for this protein is responsible for functional diversity thus producing a pseudogene. The stop codon is more frequent in African-Americans than in non-Africans.,similarity:Belongs to the G-protein coupled receptor 1 family.,
Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated
	Naniing PVahasiansa tashnalagy Ca. Itd

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



	transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
	Products Images
	117 Western Blot analysis of various cells using Olfactory receptor 10C1 Monoclonal Antibody
	85
	48 34
OR10C1	34
	26
	19
	(kD)

Nanjing BYabscience technology Co.,Ltd