



Olfactory receptor 12D3 Monoclonal Antibody

Catalog No BYmab-13463 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name OR12D3 Protein Name Olfactory receptor 12D3 Immunogen The antiserum was produced against synthesized peptide derived from human OR12D3. AA range;267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20*C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptorsimilarity:Belongs to the G-protein coupled receptor 1 family.		
Reactivity Human;Rat;Mouse; Applications WB Gene Name OR12D3 Protein Name Olfactory receptor 12D3 Immunogen The antiserum was produced against synthesized peptide derived from human OR12D3. AA range:267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors on a drape family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors o	Catalog No	BYmab-13463
Applications WB Gene Name OR12D3 Protein Name Olfactory receptor 12D3 Immunogen The antiserum was produced against synthesized peptide derived from human OR12D3. AA range:267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor years in are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors Share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition	Isotype	IgG
Gene Name OR12D3 Protein Name Olfactory receptor 12D3 Immunogen The antiserum was produced against synthesized peptide derived from human OR12D3. AA range:267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from sin	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 12D3 Immunogen The antiserum was produced against synthesized peptide derived from human OR12D3. AA range:267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a response that triggers the perception of a smell. The olfactory receptor colactory receptor colarce a receptors (GPCR) arising from single coding-exon genes. Olfactory receptor solare a 7-transmembrane domain structure with many neurotransmi	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human OR12D3. AA range:267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmerbrane domain structure with many neuronamitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of dodrant signals. The olfactory receptor gene family is the largest in	Gene Name	OR12D3
OR12D3. AA range:267-316 Specificity Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 anily, Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Protein Name	Olfactory receptor 12D3
Olfactory receptor 12D3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors or a sing from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory gene family is the largest in	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Specificity	Olfactory receptor 12D3 Monoclonal Antibody detects endogenous levels of Olfactory receptor 12D3 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90% Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Concentration	1 mg/ml
Synonyms OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Synonyms	OR12D3; Olfactory receptor 12D3; Hs6M1-27; Olfactory receptor OR6-27
Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Observed Band	35kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Tissue Specificity	Testis,
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Function	
	Background	neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in


Nanjing BYabscience technology Co.,Ltd





	proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658