



## Olfactory receptor 3A2/3A3 Monoclonal Antibody

OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 M Olfactory receptor 3A2/3A3 prFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; Olfactory	gainst synthesized peptide derived from human noclonal Antibody detects endogenous levels of
ReactivityHuman;Rat;Mouse;ApplicationsWBGene NameOR3A2/OR3A3Protein NameOlfactory receptor 3A2/3ImmunogenThe antiserum was produced OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 priorFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OIfactory receptor 3A8; Olfactory receptor 3A8;	noclonal Antibody detects endogenous levels of
ApplicationsWBGene NameOR3A2/OR3A3Protein NameOlfactory receptor 3A2/3ImmunogenThe antiserum was produced OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 M Olfactory receptor 3A2/3A3 piFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; Olfactory receptor 3A8; Olfactory receptor 3A8; Olfactory receptor 3A2; Olfactory receptor 3A8; Olfactory receptor	noclonal Antibody detects endogenous levels of
Gene NameOR3A2/OR3A3Protein NameOlfactory receptor 3A2/3ImmunogenThe antiserum was produced OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 M Olfactory receptor 3A2/3A3 ptFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; Olfactory receptor 3A2; Olfactory receptor 3A3; Olfactory receptor 3A2; Olfactory recept	noclonal Antibody detects endogenous levels of
Protein NameOlfactory receptor 3A2/3ImmunogenThe antiserum was produced OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 M Olfactory receptor 3A2/3A3 primeFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; Olfactory receptor 3A8; Olfactory receptor 3A2; Olfactory receptor 3A8; Olfactory receptor 3A2; Olfactory receptor 3A8; Olfactory receptor 3A8; Olfactory receptor 3A8; Olfactory receptor 3A8; Olfactory receptor 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass m	noclonal Antibody detects endogenous levels of
ImmunogenThe antiserum was produced OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 M Olfactory receptor 3A2/3A3 piFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; Olfactory receptor 3A8; Olfactory receptor 3A2; Olfactory receptor 3A8; Olfactory receptor 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass m	noclonal Antibody detects endogenous levels of
OR3A2/3. AA range:65-114SpecificityOlfactory receptor 3A2/3A3 M Olfactory receptor 3A2/3A3 plFormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass m	noclonal Antibody detects endogenous levels of
FormulationLiquid in PBS containing 50%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory receptor 3A8; Olfactory recep	noclonal Antibody detects endogenous levels of
SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OIfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass meta	itein.
PurificationThe antibody was affinity-puri affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass methods	glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography usingDilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory receptor 3A8;	
Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass methods	•
Purity≥90%Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass methods	
Storage Stability-20°C/1 yearSynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass methods	
SynonymsOR3A3; OR3A6; OR3A7; OR 17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass methods	
17-201; OR17-201; Olfactory receptor 3A8; Olfactory recep 3A2; OlfactoryObserved Band36kDCell PathwayCell membrane; Multi-pass methods	
Cell Pathway Cell membrane; Multi-pass m	A8P; Olfactory receptor 3A3; Olfactory receptor
	eceptor 3A6; Olfactory receptor 3A7; Olfactory or OR17-22; OR3A2; OLFRA04; Olfactory receptor
Tissue Specificity	eceptor 3A6; Olfactory receptor 3A7; Olfactory
	eceptor 3A6; Olfactory receptor 3A7; Olfactory or OR17-22; OR3A2; OLFRA04; Olfactory receptor
Function caution:It is uncertain whether receptor .,similarity:Belongs to	eceptor 3A6; Olfactory receptor 3A7; Olfactory or OR17-22; OR3A2; OLFRA04; Olfactory receptor
Background olfactory receptor family 3 sub Olfactory receptors interact win neuronal response that trigger proteins are members of a lar	eceptor 3A6; Olfactory receptor 3A7; Olfactory or OR17-22; OR3A2; OLFRA04; Olfactory receptor

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

ШR 精准的 WB 检测服务 24H 在线服务,欢迎咨询

	arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images				
	117	Western Blot analysis of various cells using Olfactory receptor 3A2/3A3 Monoclonal Antibody		
	85			
	48			
OR3A2/3A3	34			
	26			
	19 (kD)			

Nanjing BYabscience technology Co.,Ltd