



## T2R39 Monoclonal Antibody

Catalog NoBYmab-13683IsotypeIgGReactivityHuman;Rat;Mouse;ApplicationsWBGene NameTAS2R39Protein NameTaste receptor type 2 member 39ImmunogenThe antiserum was produced against synthesized peptide derived from human TAS2R39. AA range:218-267SpecificityT2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse.IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity290%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayKempsead in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.FunctionEyressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.FunctionEyressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.FunctionEyressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Revts., Sinselaneoux, Most taste cells can discriminate appna gustducin-jonkity dust taste cells can discriminate appna gustducin-jonkity dust taste cells can discriminate appna gustducin-jonkity dust taste cells can discriminate ap		
ReactivityHuman;Rat;Mouse;ApplicationsWBGene NameTAS2R39Protein NameTaste receptor type 2 member 39ImmunogenThe antiserum was produced against synthesized peptide derived from human TAS2R39. AA range:218-267SpecificityT2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Functionfunction,Receptor that may play a role in the perception of bitteress and is gustouin-positive cells.Functionfunction,Receptor that may play a role in the perception of bitteress and is gustouin-positive cells.Functionfunction,Receptor that may play a role in the perception of bitteress and is gustouin-positive cells.Functionfunction,Receptor that may play a role in the perceptor of bitteress and is gustouin-positive cells.Gene the schwade and is subsets of taste receptor reals of the tongue and exclusively in gustouin-positive cells.Gene the schwade and is gustouin-positive cells.function TReceptor that may play a role in the perceptor on posit	Catalog No	BYmab-13683
ApplicationsWBGene NameTAS2R39Protein NameTaste receptor type 2 member 39ImmunogenThe antiserum was produced against synthesized peptide derived from human TAS2R39. AA range:218-267SpecificityT2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction: Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor ray stimulate alpha gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor ray stimulate alpha gustducin, mediate PLC-beta-2 culvation and lead to the gasting of TRPM5, mixeellaneous:Most taste cells can be obscription cells of the tongue and secting of TRPM5, mixeellaneous to taste receptor cells of the tongue and beam gustducin, mediate PLC-beta-2 culvation and lead to the gas	Isotype	lgG
Gene NameTAS2R39Protein NameTaste receptor type 2 member 39ImmunogenThe antiserum was produced against synthesized peptide derived from human TAS2R39. AA range:218-267SpecificityT2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction: Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor ray stimulate alpha gustducin, mediate PLC-beta-2 cutvation and lead to the gaing of TRPM5, miscellaneous:Most taste cells can be activity of the tongue and servering of TRPM5, miscellaneous:Most taste cells can be activity of the tongue and servering of TRPM5, miscellaneous:Most taste cells can be carbon cells of the tongue and servering tasto in subset of taste receptor cells of the tongue and servering tasto in the perception of bitter composition of bitter servering tasto in servering ta	Reactivity	Human;Rat;Mouse;
Protein NameTaste receptor type 2 member 39ImmunogenThe antiserum was produced against synthesized peptide derived from human TAS2R39. AA range:218-267SpecificityT2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in guistiduin-positive cells.Functionfunction; Receptor that may play a role in the perception of bitterness and is guistiduin, mediate PLC-beta-2 activation and lead to the gain of the gastrointestinal content. The activity of this receptor Tag family, tissue specificity: Expressed in subsets of taste receptor cells of the tongue and exclusively in guistiduin, mediate PLC-beta-2 activation and lead to the gain of TRPMS, miscellaneous. Most taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor TZR family, tissue specificity: Expressed in subsets of taste receptor cells of the tongue and stimule and phate guistiducin, mediate cells can discriminate and phate guistiducin, mediate cells can discriminate and phate guistiducin, media	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human TAS2R39. AA range:218-267   Specificity T2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.   Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms TAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57   Observed Band 36kD   Cell Pathway Membrane; Multi-pass membrane protein.   Tissue Specificity Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.   Function function:Receptor that may play a role in the perception of bitterness and is gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5, miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor T2R family., tissue specificity:Expressed in subsets of taste receptor cells of the tongue and security of this receptor real sof the tongue and security of bitterexpressed in subsets	Gene Name	TAS2R39
TAS2R39. AA range:218-267SpecificityT2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor my stimulate alpha gustducin, smiscellaneous:Most taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor r2R family, tissue specificity:Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin, function; Receptor may bay a role in coupled receptor r2R family, tissue specificity:Expressed in subsets of taste receptor cells of the tongue and pilter stimuli, similarity:Belongs to the G-protein coupled receptor r2R family, tissue specificity:Expressed in subsets of taste receptor cells of the tongue and specificity:Expressed in subsets of taste receptor cells of the tongue and specificity:Expressed in subsets of ta	Protein Name	Taste receptor type 2 member 39
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms TAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57   Observed Band 36kD   Cell Pathway Membrane; Multi-pass membrane protein.   Tissue Specificity Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.   Function function:Receptor that may play a role in the perception and lead to the gating of TRPMS, miscellaneous; Most taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor T2R family, tissue specificity: cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein cells of the tongue and exclusiver of bitter compounds; individual taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor T2R family, tissue specificity: Expressed in subsets of taste receptor cells of the tongue and end of the gating of TRPMS.	Immunogen	
SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of THPM5, miscellaneoux!Most faste receptor cells of the tongue and exclusively of bitter compounds; individual taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli, similarity. Belongs to taste receptor cells of the tongue and gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5, miscellaneoux!Most faste receptor receptor T2R family, tissue sepecificity:	Specificity	T2R39 Monoclonal Antibody detects endogenous levels of T2R39 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms TAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57   Observed Band 36kD   Cell Pathway Membrane; Multi-pass membrane protein.   Tissue Specificity Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.   Function function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5, miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among biter stimuli, similarity. Evenessed in subset of taste receptor cells of the tongue and set of bitter compounds; individual taste cells can discriminate among biter stimuli, similarity. Evenessed in subset of taste receptor cells of the tongue and set of bitter compounds; individual taste cells can discriminate among biter stimuli, similarity. Evenessed in subset of taste receptor cells of the tongue and set of the tongue and set of the taste cells can discriminate among biter stimuli, similarity. Evenessed in subset of taste receptor cells of the tongue and set of the taste cells can discrim	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-inked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin-linked state cells can discriminate among bitter stimuli.similarity:Belongs to the G-protein coupled receptor T2R family.tissue stimuli.similarity:Belongs to the to captote receptor cells of the tongue and stimuli.similarity:Belongs to the state receptor cells of the tongue and stimuli.similarity:Belongs to the state receptor cells of the tongue and stimuli.similarity:Belongs to the G-protein coupled receptor T2	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms TAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57   Observed Band 36kD   Cell Pathway Membrane; Multi-pass membrane protein.   Tissue Specificity Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.   Function function:Receptor that may play a role in the perception of bitterness and is gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPMS, miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor T2R family, tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Purification	
Purity ≥90%   Storage Stability -20°C/1 year   Synonyms TAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57   Observed Band 36kD   Cell Pathway Membrane; Multi-pass membrane protein.   Tissue Specificity Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.   Function function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5, miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli, similarity:Belongs to the G-protein coupled receptor T2R family., tissue specificity: Expressed in subsets of taste receptor cells of the tongue and	Dilution	WB 1:500-2000
Storage Stability-20°C/1 yearSynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Concentration	1 mg/ml
SynonymsTAS2R39; Taste receptor type 2 member 39; T2R39; Taste receptor type 2 member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Purity	≥90%
member 57; T2R57Observed Band36kDCell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Storage Stability	-20°C/1 year
Cell PathwayMembrane; Multi-pass membrane protein.Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Synonyms	
Tissue SpecificityExpressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.Functionfunction:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Observed Band	36kD
Function function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Cell Pathway	Membrane; Multi-pass membrane protein.
gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and	Tissue Specificity	Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.
	Function	gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商


精准的 WB 检测服务

24H 在线服务,欢迎咨询



Background	The protein encoded by this gene is a bitter taste receptor that detects green tea catechins, soy isoflavones, and theaflavins. The encoded protein is gustducin-linked and may activate alpha gustducin. This gene is intronless. [provided by RefSeq, Dec 2015],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd