



T2R7 Monoclonal Antibody

Catalog No	BYmab-13689
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TAS2R7
Protein Name	Taste receptor type 2 member 7
Immunogen	The antiserum was produced against synthesized peptide derived from human TAS2R7. AA range:47-96
Specificity	T2R7 Monoclonal Antibody detects endogenous levels of T2R7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TAS2R7; Taste receptor type 2 member 7; T2R7; Taste receptor family B member 4; TRB4
Observed Band	36kD
Cell Pathway	Membrane; Multi-pass membrane protein.
Tissue Specificity	Expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells.
Function	function:Gustducin-coupled receptor implicated in the perception of bitter compounds in the oral cavity and the gastrointestinal tract. Signals through PLCB2 and the calcium-regulated cation channel TRPM5.,miscellaneous:Several bitter taste receptors are expressed in a single taste receptor cell.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells.,
Background	This gene product belongs to the family of candidate taste receptors that are members of the G-protein-coupled receptor superfamily. These proteins are
	Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务, 欢迎咨询



	specifically expressed in the taste receptor cells of the tongue and palate epithelia. They are organized in the genome in clusters and are genetically linked to loci that influence bitter perception in mice and humans. In functional expression studies, they respond to bitter tastants. This gene maps to the taste receptor gene cluster on chromosome 12p13. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Nanjing BYabscience technology Co.,Ltd