



IL-1RI Monoclonal Antibody

Catalog No	BYmab-13720
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	IL1R1
Protein Name	Interleukin-1 receptor type 1
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human IL1R1. AA range:31-80
Specificity	IL-1RI Monoclonal Antibody detects endogenous levels of IL-1RI protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IL1R1; IL1R; IL1RA; IL1RT1; Interleukin-1 receptor type 1; IL-1R-1; IL-1RT-1; IL-1RT1; CD121 antigen-like family member A; Interleukin-1 receptor alpha; IL-1R-alpha; Interleukin-1 receptor type I; p80; CD121a
Observed Band	75kD
Cell Pathway	Membrane ; Single-pass type I membrane protein . Cell membrane . Secreted .
Tissue Specificity	Expressed in T-helper cell subsets. Preferentially expressed in T-helper 1 (Th1) cells.
Function	function:Receptor for interleukin-1 alpha (IL-1A), beta (IL-1B), and interleukin-1 receptor antagonist protein (IL-1RA). Binding to the agonist leads to the activation of NF-kappa-B. Signaling involves formation of a ternary complex containing IL1RAP, TOLLIP, MYD88, and IRAK1 or IRAK2.,similarity:Belongs to the interleukin-1 receptor family.,similarity:Contains 1 TIR domain.,similarity:Contains 3 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Binds IL1RAP.,
Background	This gene encodes a cytokine receptor that belongs to the interleukin-1 receptor family. The encoded protein is a receptor for interleukin-1 alpha, interleukin-1
	Naniing BYabscience technology Co. Ltd

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务, 欢迎咨询

beta, and interleukin-1 receptor antagonist. It is an important mediator involved in many cytokine-induced immune and inflammatory responses. This gene is located in a cluster of related cytokine receptor genes on chromosome 2q12. [provided by RefSeq, Dec 2013], Avoid repeated freezing and thawing! matters needing attention This product can be used in immunological reaction related experiments. For Usage suggestions more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IL-1RI Monoclonal Antibody

Nanjing BYabscience technology Co., Ltd