




SAMHD1 (phospho-Thr592) mouse mAb

Catalog No	BYmab-13861
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SAMHD1 MOP5
Protein Name	SAMHD1 (Thr592)
Immunogen	Synthesized phosho peptide around human SAMHD1 (Thr592)
Specificity	This antibody detects endogenous levels of Human Mouse Rat SAMHD1 (phospho-Thr592)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
	. 000/
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	-20°C/1 year SAM domain and HD domain-containing protein 1 (EC 3.1.4) (Dendritic
Storage Stability Synonyms	-20°C/1 year SAM domain and HD domain-containing protein 1 (EC 3.1.4) (Dendritic cell-derived IFNG-induced protein) (DCIP) (Monocyte protein 5) (MOP-5)
Storage Stability Synonyms Observed Band	-20°C/1 year SAM domain and HD domain-containing protein 1 (EC 3.1.4) (Dendritic cell-derived IFNG-induced protein) (DCIP) (Monocyte protein 5) (MOP-5) 72kD Nucleus . Chromosome . Localizes to sites of DNA double-strand breaks in
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year SAM domain and HD domain-containing protein 1 (EC 3.1.4) (Dendritic cell-derived IFNG-induced protein) (DCIP) (Monocyte protein 5) (MOP-5) 72kD Nucleus . Chromosome . Localizes to sites of DNA double-strand breaks in response to DNA damage Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes (PubMed:11064105). No expression is seen in
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year SAM domain and HD domain-containing protein 1 (EC 3.1.4) (Dendritic cell-derived IFNG-induced protein) (DCIP) (Monocyte protein 5) (MOP-5) 72kD Nucleus . Chromosome . Localizes to sites of DNA double-strand breaks in response to DNA damage Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes (PubMed:11064105). No expression is seen in brain and thymus (PubMed:11064105). function:May play a role in mediating proinflammatory responses to TNF-alpha signaling.,induction:By interferon gamma. Up-regulated in TNF-alpha treated lung fibroblasts.,similarity:Contains 1 HD domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,tissue specificity:Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes. No

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







response to viral infection and may be involved in mediation of tumor necrosis factor-alpha proinflammatory responses. Mutations in this gene have been associated with Aicardi-Goutieres syndrome. [provided by RefSeq, Mar 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using SAMHD1 (phospho-Thr592) mouse mAb

Nanjing BYabscience technology Co.,Ltd