



# C9 Monoclonal Antibody

<b>Catalog No</b>	BYmab-13877
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	C9
<b>Protein Name</b>	Complement component C9
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human C9. AA range:181-230
<b>Specificity</b>	C9 Monoclonal Antibody detects endogenous levels of C9 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	C9; Complement component C9
<b>Observed Band</b>	70kD
<b>Cell Pathway</b>	Secreted . Target cell membrane ; Multi-pass membrane protein . Secreted as soluble monomer. Oligomerizes at target membranes, forming a pre-pore. A conformation change then leads to the formation of a 100 Angstrom diameter pore. .
<b>Tissue Specificity</b>	Plasma (at protein level).
<b>Function</b>	disease:Defects in C9 are a cause of component C9 deficiency (C9D) [MIM:120940]. Patients with C9D suffer from recurrent bacterial infections, predominantly from Neisseria meningitidis.,function:C9 is the final component of the complement system to be added in the assembly of the membrane attack complex. It is able to enter lipid bilayers, forming transmembrane channels.,online information:C9 mutation db,PTM:Thrombin cleaves factor C9 to produce C9a and C9b.,similarity:Belongs to the complement C6/C7/C8/C9 family.,similarity:Contains 1 EGF-like domain.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 1 MACPF domain.,similarity:Contains 1 TSP

**Nanjing BYabscience technology Co.,Ltd**



type-1 domain.,

#### Background

This gene encodes the final component of the complement system. It participates in the formation of the Membrane Attack Complex (MAC). The MAC assembles on bacterial membranes to form a pore, permitting disruption of bacterial membrane organization. Mutations in this gene cause component C9 deficiency. [provided by RefSeq, Feb 2009],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Western Blot analysis of various cells using C9 Monoclonal Antibody

**Nanjing BYabs science technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658