



CD314 Monoclonal Antibody

Catalog No	BYmab-13895
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	KLRK1
Protein Name	NKG2-D type II integral membrane protein
Immunogen	The antiserum was produced against synthesized peptide derived from human KLRK1. AA range:111-160
Specificity	CD314 Monoclonal Antibody detects endogenous levels of CD314 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KLRK1; D12S2489E; NKG2D; NKG2-D type II integral membrane protein; Killer cell lectin-like receptor subfamily K member 1; NK cell receptor D; NKG2-D-activating NK receptor; CD antigen CD314
Observed Band	25kD
Cell Pathway	Cell membrane ; Single-pass type II membrane protein . Colocalized with HCST on the cell surface.
Tissue Specificity	Expressed in natural killer (NK) cells, CD8(+) alpha-beta and gamma-delta T-cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Expressed in interferon-producing killer dendritic cells (IKDCs).
Function	alternative products:A number of isoforms are produced,function:Receptor for MICA, MICB, ULBP1, ULBP2, ULBP3 (ULBP2>ULBP1>ULBP3) and ULBP4. Plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells. Involved in the immune surveillance exerted by T- and B-lymphocytes.,miscellaneous:Structurally distinct families of ligands for mouse and human NKG2D receptors have been characterized. They might be orthologs.,online information:NKG-2D,similarity:Contains 1 C-type lectin

Nanjing BYabscience technology Co.,Ltd



domain.,subunit:Homodimer. Interacts with DAP10. The interaction with DAP10 is required for NKG2D cell surface expression.,tissue specificity:Natural killer cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Also detected in gamma-delta cells and CD8+ alpha-beta T-cells. Expressed in interferon-producing killer dendritic cells (IKDCs).

Background

Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD314 Monoclonal Antibody