



CD32-C Monoclonal Antibody


| | |
|---------------------------|--|
| Catalog No | BYmab-13898 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | FCGR2C |
| Protein Name | Low affinity immunoglobulin gamma Fc region receptor II-c |
| Immunogen | The antiserum was produced against synthesized peptide derived from human FCGR2C. AA range:251-300 |
| Specificity | CD32-C Monoclonal Antibody detects endogenous levels of CD32-C protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | FCGR2C; CD32; FCG2; IGFR2; Low affinity immunoglobulin gamma Fc region receptor II-c; IgG Fc receptor II-c; CDw32; Fc-gamma RII-c; Fc-gamma-RIIc; FcRII-c; CD antigen CD32 |
| Observed Band | 35kD |
| Cell Pathway | [Isoform IIC4]: Cytoplasm .; [Isoform IIC3]: Cell membrane; Single-pass type I membrane protein.; [Isoform IIC2]: Cell membrane; Single-pass type I membrane protein.; [Isoform IIC1]: Cell membrane; Single-pass type I membrane protein. |
| Tissue Specificity | Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells. |
| Function | caution:Has sometimes been attributed to correspond to FcR-IIb.,caution:Has sometimes been attributed to correspond to FcR-IIC.,disease:A chromosomal aberration involving FCGR2B is found in a follicular lymphoma. Translocation t(1;22)(q22;q11). The translocation leads to the hyperexpression of the receptor. This may play a role in the tumor progression.,domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. |

Nanjing BYabscience technology Co.,Ltd



| | |
|---|---|
| <p>Background</p> | <p>The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,domain:Contains an intracytoplasmic twice repeated motif referred as immunoreceptor tyrosine-based activator motif (ITAM). These motifs are involved in triggering cell activation upon receptors aggregation.,function:Receptor for the Fc region of complexed immunoglobulin gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.,function:Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Isoform IIB1 interacts with measles virus N protein. N protein is released in the blood following lysis of measles infected cells. This interaction presumably block inflammatory immune response. Interacts with INPP5D/SHIP1.,tissue specificity:Is the most broadly distributed Fc-gamma-receptor. Expressed in monocyte, neutrophils, macrophages, basophils, eosinophils, Langerhans cells, B-cells, platelets cells and placenta (endothelial cells). Not detected in natural killer cells.,tissue specificity:Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells.,</p> |
| <p>matters needing attention</p> | <p>Avoid repeated freezing and thawing!</p> |
| <p>Usage suggestions</p> | <p>This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.</p> |

Products Images



Western Blot analysis of various cells using CD32-C Monoclonal Antibody

nce technology Co.,Ltd

025-5229-8998

监督电话: 15950492658