



CD1D Monoclonal Antibody


| | |
|---------------------------|---|
| Catalog No | BYmab-13980 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | CD1D |
| Protein Name | Antigen-presenting glycoprotein CD1d |
| Immunogen | Synthesized peptide derived from Antigen-presenting glycoprotein CD1d at AA range: 161-210 |
| Specificity | CD1D Monoclonal Antibody detects endogenous levels of CD1D protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | CD1D; Antigen-presenting glycoprotein CD1d; R3G1; CD1d |
| Observed Band | 37kD |
| Cell Pathway | Cell membrane ; Single-pass type I membrane protein . Basolateral cell membrane ; Single-pass type I membrane protein . Endosome membrane ; Single-pass type I membrane protein . Lysosome membrane ; Single-pass type I membrane protein . Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Subject to intracellular trafficking between the cell membrane, endosomes and lysosomes. . |
| Tissue Specificity | Expressed on cortical thymocytes, on certain T-cell leukemias, and in various other tissues. |
| Function | function:Antigen-presenting protein that binds self and non-self glycolipids and presents them to T-cell receptors on natural killer T-cells.,miscellaneous:During protein synthesis and maturation, CD1 family members bind endogenous lipids that are replaced by lipid or glycolipid antigens when the proteins are internalized and pass through endosomes, before trafficking back to the cell surface.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain.,subcellular location:Subject to intracellular trafficking between the cell membrane, |

Nanjing BYabscience technology Co.,Ltd



| | |
|----------------------------------|--|
| Background | <p>endosomes and lysosomes.,subunit:Heterodimer with B2M (beta-2-microglobulin). Interacts with MHC II.,tissue specificity:Expressed on cortical thymocytes, on certain T-cell leukemias, and in various other tissues.,</p> <p>This gene encodes a divergent member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to late endosomes and lysosomes via a tyrosine-based motif in the cytoplasmic tail. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2016],</p> |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Western Blot analysis of various cells using CD1D Monoclonal Antibody