



SAMHD1 Monoclonal Antibody

Catalog No	BYmab-14039
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	SAMHD1
Protein Name	SAM domain and HD domain-containing protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human SAMHD1. AA range:431-480
Specificity	SAMHD1 Monoclonal Antibody detects endogenous levels of SAMHD1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
	•
Synonyms	SAMHD1; MOP5; SAM domain and HD domain-containing protein 1; Dendritic cell-derived IFNG-induced protein; DCIP; Monocyte protein 5; MOP-5
Synonyms Observed Band	SAMHD1; MOP5; SAM domain and HD domain-containing protein 1; Dendritic
	SAMHD1; MOP5; SAM domain and HD domain-containing protein 1; Dendritic cell-derived IFNG-induced protein; DCIP; Monocyte protein 5; MOP-5
Observed Band	SAMHD1; MOP5; SAM domain and HD domain-containing protein 1; Dendritic cell-derived IFNG-induced protein; DCIP; Monocyte protein 5; MOP-5 72kD Nucleus . Chromosome . Localizes to sites of DNA double-strand breaks in
Observed Band Cell Pathway	SAMHD1; MOP5; SAM domain and HD domain-containing protein 1; Dendritic cell-derived IFNG-induced protein; DCIP; Monocyte protein 5; MOP-5 72kD Nucleus . Chromosome . Localizes to sites of DNA double-strand breaks in response to DNA damage Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes (PubMed:11064105). No expression is seen in brain and thymus (PubMed:11064105). function:May play a role in mediating proinflammatory responses to TNF-alpha
Observed Band Cell Pathway Tissue Specificity	SAMHD1; MOP5; SAM domain and HD domain-containing protein 1; Dendritic cell-derived IFNG-induced protein; DCIP; Monocyte protein 5; MOP-5 72kD Nucleus . Chromosome . Localizes to sites of DNA double-strand breaks in response to DNA damage Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes (PubMed:11064105). No expression is seen in brain and thymus (PubMed:11064105). function:May play a role in mediating proinflammatory responses to TNF-alpha signaling.,induction:By interferon gamma. Up-regulated in TNF-alpha treated lung fibroblasts.,similarity:Contains 1 HD domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,tissue specificity:Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes. No

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	regulation of the innate immune response. The encoded protein is upregulated in response to viral infection and may be involved in mediation of tumor necrosis factor-alpha proinflammatory responses. Mutations in this gene have been associated with Aicardi-Goutieres syndrome. [provided by RefSeq, Mar 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Western Blot analysis of various cells using SAMHD1 Monoclonal Antibody Western Blot analysis of various cells using SAMHD1 Monoclonal Antibody

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658