



# AID Monoclonal Antibody

<b>Catalog No</b>	BYmab-14045
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	AICDA
<b>Protein Name</b>	Activation-induced cytidine deaminase
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human AICDA. AA range:81-130
<b>Specificity</b>	AID Monoclonal Antibody detects endogenous levels of AID protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	AICDA; AID; Activation-induced cytidine deaminase; Cytidine aminohydrolase
<b>Observed Band</b>	24kD
<b>Cell Pathway</b>	Nucleus . Cytoplasm . Predominantly cytoplasmic (PubMed:21385873). In the presence of MCM3AP/GANP, relocates to the nucleus (By similarity). .
<b>Tissue Specificity</b>	Strongly expressed in lymph nodes and tonsils.
<b>Function</b>	catalytic activity:Cytidine + H(2)O = uridine + NH(3).,cofactor:Zinc.,disease:Defects in AICDA are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2) [MIM:605258]. HIGM2 is characterized by normal or elevated serum IgM levels with absence of IgG, IgA, and IgE, resulting in a profound susceptibility to bacterial infections. HIGM2 causes the absence of Ig class switch recombination (CSR), the lack of Ig somatic hypermutations, and lymph node hyperplasia caused by the presence of giant germinal centers.,function:RNA-editing deaminase involved in somatic hypermutation, gene conversion, and class-switch recombination. Required for several crucial steps of B-cell terminal differentiation necessary for efficient antibody responses.,online information:AICDA mutation db,similarity:Belongs to


**Nanjing BYabscience technology Co.,Ltd**



the cytidine and deoxycytidylate deaminase family.,tissue specificity:Str

<b>Background</b>	This gene encodes a RNA-editing deaminase that is a member of the cytidine deaminase family. The protein is involved in somatic hypermutation, gene conversion, and class-switch recombination of immunoglobulin genes. Defects in this gene are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2). [provided by RefSeq, Feb 2009],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using AID Monoclonal Antibody

**Nanjing BYabs science technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658