



Pim-1 (phospho Tyr309) Monoclonal Antibody

Catalog No	BYmab-14368
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PIM1
Protein Name	Serine/threonine-protein kinase pim-1
Immunogen	The antiserum was produced against synthesized peptide derived from human Pim-1 around the phosphorylation site of Tyr309. AA range:281-330
Specificity	Phospho-Pim-1 (Y309) Monoclonal Antibody detects endogenous levels of Pim-1 protein only when phosphorylated at Y309.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PIM1; Serine/threonine-protein kinase pim-1
Observed Band	45kD
Cell Pathway	[Isoform 1]: Cytoplasm. Nucleus.; [Isoform 2]: Cell membrane.
Tissue Specificity	Expressed primarily in cells of the hematopoietic and germline lineages. Isoform 1 and isoform 2 are both expressed in prostate cancer cell lines.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.;cofactor:Manganese.;function:Plays a role in signal transduction in blood cells. Contributes to both cell proliferation and survival and thus provide a selective advantage in tumorigenesis. May affect the structure or silencing of chromatin by phosphorylating HP1 gamma/CBX3.;induction:Strongly induced in leukocytes by the JAK/STAT pathway in response to cytokines.;PTM:Autophosphorylated on both serine/threonine and tyrosine residues.;similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. PIM subfamily.;similarity:Contains 1 protein kinase domain.;subunit:Binds to RP9. Isoform 2 is isolated as a monomer whereas isoform 1 complexes with other proteins. Isoform 1, but not isoform 2, binds

Nanjing BYabscience technology Co.,Ltd



BMX.,tissue specificity:Expressed primarily in cells of the hematopoietic and germline lineages. Isoform 1 an

Background

The protein encoded by this gene belongs to the Ser/Thr protein kinase family, and PIM subfamily. This gene is expressed primarily in B-lymphoid and myeloid cell lines, and is overexpressed in hematopoietic malignancies and in prostate cancer. It plays a role in signal transduction in blood cells, contributing to both cell proliferation and survival, and thus provides a selective advantage in tumorigenesis. Both the human and orthologous mouse genes have been reported to encode two isoforms (with preferential cellular localization) resulting from the use of alternative in-frame translation initiation codons, the upstream non-AUG (CUG) and downstream AUG codons (PMIDs:16186805, 1825810).[provided by RefSeq, Aug 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Pim-1 (phospho Tyr309) Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658