




FAK (Phospho Ser722) mouse mAb

Catalog No BYmab-14633 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name PTK2 FAK FAK1 Protein Name FAK (Phospho Ser722) Immunogen Synthesized peptide derived from human FAK (Phospho Ser722) Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; cytoskeleton, cilium basal body. Constitutent of focal adhesions. Detected at microtubule organizing center, certrosome. Nucleus. Cytoplasm, cytoskeleton inclima basal body		
Reactivity Human;Mouse;Rat Applications WB Gene Name PTK2 FAK FAK1 Protein Name FAK (Phospho Ser722) Immunogen Synthesized peptide derived from human FAK (Phospho Ser722) Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic, side. Cytoplasmic, cell cortex. Cytoplasmic, cytoskeleton. Cytoplasmic, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasmic, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasmic, cytoskeleton, piclium basal body. Constituent of focal adhesions. Detec	Catalog No	BYmab-14633
Applications Gene Name PTK2 FAK FAK1 Protein Name FAK (Phospho Ser722) Immunogen Synthesized peptide derived from human FAK (Phospho Ser722) Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic, side. Cytoplasmic, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) in differentiation, angiogenesis, blood vessel development, protein complex assembly, protein amino acid	Isotype	IgG
Gene Name PTK2 FAK FAK1 Protein Name FAK (Phospho Ser722) Immunogen Synthesized peptide derived from human FAK (Phospho Ser722) Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK,pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmi, cell cortex. Cytoplasmic, cytoskeleton. Cytoplasmi, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithe	Reactivity	Human;Mouse;Rat
Protein Name FAK (Phospho Ser722) Immunogen Synthesized peptide derived from human FAK (Phospho Ser722) Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, illum basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function	Applications	WB
Immunogen Synthesized peptide derived from human FAK (Phospho Ser722) Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasm; cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, illum basal body. Constituent of focal adhesions. Detected at microtubule cytoskeleton organization, cell morphogenesis, ell morphogenesis involved in differentiation, angiogenesis, blood vessel in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesi	Gene Name	PTK2 FAK FAK1
Specificity This antibody detects endogenous levels of Human,Mouse,Rat FAK (Phospho Ser722) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasm; cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes, Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function Tinde Time Time Time Time Time Time Time Tim	Protein Name	FAK (Phospho Ser722)
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm. Cytoskeleton. Cytoplasmic orenter, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, plood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Immunogen	Synthesized peptide derived from human FAK (Phospho Ser722)
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; cytoskeleton, microtubule organizing center, centrosome . Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus. Cytoplasm, cytoskeleton, cilium basal body . Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, neuron migration, vasculature development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Specificity	This antibody detects endogenous levels of Human, Mouse, Rat FAK (Phospho Ser722)
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, yasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus. Cytoplasm, cytoskeleton, cilium basal body . Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, neuron migration, vasculature development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Source	Monoclonal, Mouse,lgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, protein complex assembly, protein amino acid	Dilution	WB 1:500-2000
Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Concentration	1 mg/ml
Synonyms Focal adhesion kinase 1 (FADK 1;EC 2.7.10.2;Focal adhesion kinase-related nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell Pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Purity	≥90%
nonkinase;FRNK;Protein phosphatase 1 regulatory subunit 71;PPP1R71;Protein-tyrosine kinase 2;p125FAK;pp125FAK) Observed Band 125kD Cell pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Storage Stability	-20°C/1 year
Cell pathway Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation,angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Synonyms	nonkinase;FRNK;Protein phosphatase 1 regulatory subunit
Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cytoplasm, cytoskeleton, cilium basal body. Constituent of focal adhesions. Detected at microtubules. Tissue Specificity Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Observed Band	125kD
fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787). Function microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation, angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Cell Pathway	Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus. Cytoplasm, cytoskeleton, cilium basal body . Constituent of focal adhesions. Detected at
involved in differentiation,angiogenesis, blood vessel development, vasculogenesis, neuron migration, vasculature development, protein complex assembly, protein amino acid	Tissue Specificity	fibroblasts (at protein level). Ubiquitous. Expressed in epithelial cells (at protein
	Function	microtubule cytoskeleton organization, cell morphogenesis, cell morphogenesis involved in differentiation,angiogenesis, blood vessel

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







process, cell motion, cytoskeleton organization, microtubule-based process, cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway, transmembrane receptor protein tyrosine kinase signaling pathway, signal complex assembly, integrin-mediated signaling pathway, cell-cell signaling, synaptic transmission, axonogenesis, negative regulation of cell development, regulation of cell morphogenesis involved in differentiation, regulation of neuron projection development, phosphorylation, cell migration, transmission of nerve impulse, central nervo

Background

catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,domain:The carboxy-terminal region is the site of focal adhesion targeting (FAT) sequence which mediates the localization of FAK1 to focal adhesions.,domain:The first Pro-rich domain interacts with the SH3 domain of CRK-associated substrate (BCAR1) and CASL.,function:Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity.,PTM:Phosphorylated on 6 tyrosine residues upon activation.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family, similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family, FAK subfamily.,similarity:Contains 1 FERM domain.,similarity:Contains 1 protein kinase domain.,subcellular location:Constituent of focal adhesions.,subunit:Interacts with CAS family members and with GIT1, SORBS1 and BCAR3. Interacts with RGNEF and SHB (By similarity). Interacts with TGFB111.,tissue specificity:Expressed in all organs tested, in lymphoid cell lines, but most abundantly in brain.,

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using FAK (Phospho Ser722) mouse mAb



nce technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658