



ADK Monoclonal Antibody

Catalog No	BYmab-14647
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ADK
Protein Name	Adenosine kinase
Immunogen	The antiserum was produced against synthesized peptide derived from human ADK. AA range:1-50
Specificity	ADK Monoclonal Antibody detects endogenous levels of ADK protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ADK; Adenosine kinase; AK; Adenosine 5'-phosphotransferase
Observed Band	40kD
Cell Pathway	[Isoform 1]: Nucleus .; [Isoform 2]: Cytoplasm .
Tissue Specificity	Widely expressed. Highest level in placenta, liver, muscle and kidney.
Function	catalytic activity:ATP + adenosine = ADP + AMP.,cofactor:Binds 3 magnesium ions per subunit.,function:ATP dependent phosphorylation of adenosine and other related nucleoside analogs to monophosphate derivatives. Serves as a potential regulator of concentrations of extracellular adenosine and intracellular adenine nucleotides.,pathway:Purine metabolism; AMP biosynthesis via salvage pathway; AMP from adenosine: step 1/1.,similarity:Belongs to the carbohydrate kinase pfkB family.,subunit:Monomer.,tissue specificity:Widely expressed. Highest level in placenta, liver, muscle and kidney.,
Background	This gene an enzyme which catalyzes the transfer of the gamma-phosphate from ATP to adenosine, thereby serving as a regulator of concentrations of both

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



extracellular adenosine and intracellular adenine nucleotides. Adenosine has widespread effects on the cardiovascular, nervous, respiratory, and immune systems and inhibitors of the enzyme could play an important pharmacological role in increasing intravascular adenosine concentrations and acting as anti-inflammatory agents. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd