



Dyrk1B Monoclonal Antibody

Catalog No	BYmab-14735
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DYRK1B
Protein Name	Dual specificity tyrosine-phosphorylation-regulated kinase 1B
Immunogen	The antiserum was produced against synthesized peptide derived from human DYRK1B. AA range:331-380
Specificity	Dyrk1B Monoclonal Antibody detects endogenous levels of Dyrk1B protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DYRK1B; MIRK; Dual specificity tyrosine-phosphorylation-regulated kinase 1B; Minibrain-related kinase; Mirk protein kinase
Observed Band	70kD
Cell Pathway	Nucleus . Nucleus, nucleolus . Chromosome . Localizes to sites of double-strand breaks (DSBs) following DNA damage. .
Tissue Specificity	Highest expression in skeletal muscle, testis, heart and brain with little expression in colon or lung. Expressed in a variety of tumor cell lines.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Inhibited by RANBP9.,function:Dual-specificity kinase which possesses both serine/ threonine and tyrosine kinase activities. Enhances the transcriptional activity of TCF1/HNF1A and FOXO1. Inhibits epithelial cell migration. Mediates colon carcinoma cell survival in mitogen-poor environments.,PTM:Autophosphorylated on tyrosine residues. Phosphorylated by MAP kinase. Tyrosine phosphorylation may be required for dimerization.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MNB/DYRK subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Dimer. Interacts with DCOHM, MAP2K3/MKK3, RANBP9 and

Nanjing BYabscience technology Co.,Ltd



TCF1/HNF1A. Part of a complex consisting of RANBP9, RAN, DYRK1B and COPS5.,tissue specificity: Highest expression in skeletal muscle, testis, heart and brain with little expressio

Background

This gene encodes a member of a family of nuclear-localized protein kinases. The encoded protein participates in the regulation of the cell cycle. Expression of this gene may be altered in tumor cells, and mutations in this gene were found to cause abdominal obesity-metabolic syndrome 3. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd