



## **GRK 3 Monoclonal Antibody**

| Catalog No         | BYmab-14764   |
|--------------------|---|
| Isotype            | IgG   |
| Reactivity         | Human;Mouse;Rat   |
| Applications       | WB  |
| Gene Name          | ADRBK2  |
| Protein Name       | Beta-adrenergic receptor kinase 2   |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human GRK3. AA range:361-410  |
| Specificity        | GRK 3 Monoclonal Antibody detects endogenous levels of GRK 3 protein.   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Monoclonal, Mouse,IgG   |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution           | WB 1:500-2000   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | ADRBK2; BARK2; GRK3; Beta-adrenergic receptor kinase 2; Beta-ARK-2;<br>G-protein-coupled receptor kinase 3  |
| Observed Band      | 80kD  |
| Cell Pathway       | Cell junction, synapse, postsynapse . Cell junction, synapse, presynapse .  |
| Tissue Specificity | Brain,Ovary,  |
| Function           | catalytic activity:ATP + [beta-adrenergic receptor] = ADP + [beta-adrenergic receptor] phosphate.,function:Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily., GPRK subfamily.,similarity:Contains 1 AGC Ser/Thr protein kinase family. GPRK subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 RGS domain., |
|                    |   |

## Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务,欢迎咨询



| Background                | The beta-adrenergic receptor kinase specifically phosphorylates the agonist-occupied form of the beta-adrenergic and related G protein-coupled receptors. Overall, the beta adrenergic receptor kinase 2 has 85% amino acid similarity with beta adrenergic receptor kinase 1, with the protein kinase catalytic domain having 95% similarity. These data suggest the existence of a family of receptor kinases which may serve broadly to regulate receptor function. [provided by RefSeq, Jul 2008], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing!   |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |
|                           |  |

## **Products Images**



Western Blot analysis of various cells using GRK 3 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd