



TTK Monoclonal Antibody

Catalog No	BYmab-15024			
lsotype	lgG			
Reactivity	Human;Mouse			
Applications	WB			
Gene Name	ТТК			
Protein Name	Dual specificity protein kinase TTK			
Immunogen	The antiserum was produced against synthesized peptide derived from human TTK. AA range:642-691			
Specificity	TTK Monoclonal Antibody detects endogenous levels of TTK protein.			
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.			
Source	Monoclonal, Mouse,IgG			
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.			
Dilution	WB 1:500-2000			
Concentration	1 mg/ml			
Purity	≥90%			
Purity Storage Stability	≥90% -20°C/1 year			
-				
Storage Stability	-20°C/1 year TTK; MPS1; MPS1L1; Dual specificity protein kinase TTK; Phosphotyrosine			
Storage Stability Synonyms	-20°C/1 year TTK; MPS1; MPS1L1; Dual specificity protein kinase TTK; Phosphotyrosine picked threonine-protein kinase; PYT			
Storage Stability Synonyms Observed Band	-20°C/1 year TTK; MPS1; MPS1L1; Dual specificity protein kinase TTK; Phosphotyrosine picked threonine-protein kinase; PYT 100kD			
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year TTK; MPS1; MPS1L1; Dual specificity protein kinase TTK; Phosphotyrosine picked threonine-protein kinase; PYT 100kD kinetochore,cytoplasm,spindle,membrane,			
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	 -20°C/1 year TTK; MPS1; MPS1L1; Dual specificity protein kinase TTK; Phosphotyrosine picked threonine-protein kinase; PYT 100kD kinetochore,cytoplasm,spindle,membrane, Present in rapidly proliferating cell lines. catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Phosphorylates proteins on serine, threonine, and tyrosine. Probably associated with cell proliferation.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase domain.,tissue 			



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



	duplication. It has been found to be a critical mitotic checkpoint protein for accurate segregation of chromosomes during mitosis. Tumorigenesis may occur when this protein fails to degrade and produces excess centrosomes resulting in aberrant mitotic spindles. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2009],			
matters needing attention	Avoid repeated freezing and thawing!			
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.			
	Proc	lucts Images		
HuvEC HuvEC		Western Blot ana Monoclonal Antib	lysis of various cells using TTK ody	
ттк	117	7		
	85			
	48			
	34			
	26			
	19 (kD)			
Nanjing BYabscience technology Co.,Ltd				
网址:www.njbybio.com 官方打		: 025-5229-8998	监督电话: 15950492658	