




## MOX-2 Monoclonal Antibody

Catalog No         BYmab-15778           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         MEOX2           Protein Name         Homeobox protein MOX-2           Immunogen         The antiserum was produced against synthesized peptide derived from human MEOX2. AN range:151-200           Specificity         MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         290%           Storage Stability         -20°C/1 year           Synonyms         MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2           Observed Band         33kD           Cell Pathway         Nucleus . Nucleus speckle .           Tissue Specificity         Embryo,Heart,Skin,           Function         Embryo,Heart,Skin,           Function         Investion:Role in mesoderm induction and its earliest		
Reactivity Human;Mouse;Rat  Applications WB  Gene Name MEOX2  Protein Name Homeobox protein MOX-2  Immunogen The antiserum was produced against synthesized peptide derived from human MEOX2. AA range:151-200  Specificity MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo,Heart,Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle, polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population, similarity. Contains† homeobox DNA-binding domain. subunit.Interacts with RNF10, its sue specificity:Embryo and placenta.  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play and antennapedia-like homeobox-containing genes. The encoded protein may play a	Catalog No	BYmab-15778
Applications WB Gene Name MEOX2 Protein Name Homeobox protein MOX-2  Immunogen The antiserum was produced against synthesized peptide derived from human MEOX2. AA range:151-200 Specificity MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, lgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle, polymorphism. The poly-His region of MEOX2 is polymorphic and the nomeobox DNA-binding domain, subunit.Interacts with RNF10, issue specificity:Embryo and placenta.  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded privater may play a antennapedia-like homeobox-containing genes. The encoded privater may play a	Isotype	IgG
Gene Name MEOX2  Protein Name Homeobox protein MOX-2  Immunogen The antiserum was produced against synthesized peptide derived from human MEOX2. AA range:151-200  Specificity MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  function: Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and solerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism: The poly-His region of MEOX2 is polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population.; similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity:Embryo and placenta.  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Reactivity	Human;Mouse;Rat
Protein Name	Applications	WB
Immunogen         The antiserum was produced against synthesized peptide derived from human MEOX2. AA range:151-200           Specificity         MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         MEOX2, GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2           Observed Band         33kD           Cell Pathway         Nucleus . Nucleus speckle .           Tissue Specificity         Embryo,Heart,Skin,           Function         function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle_polymorphis and the number of His varies in the population, similarity.Contains 1 homeobox DNA-binding domain, subunit:Interacts with RNF10, tissue specificity:Embryo and placenta.           Background         This gene encodes a member of a subfamily of non-clustered, diverged, antennagedia-like homeobox-containi	Gene Name	MEOX2
MEOX2. AA range:151-200  Specificity MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle, polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population, similarity: Contains 1 homeobox DNA-binding domain, subunit:Interacts with RNF10., tissue specificity:Embryo and placenta.,  Background The pencoded protein may play a antennapedia-like homeobox-containing genes. The encoded protein may play a	Protein Name	Homeobox protein MOX-2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population, similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity: Embryo and placenta.,  Background The poly-Lise and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle, polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population, similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity: Embryo and placenta.,	lmmunogen	
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2         Observed Band       33kD         Cell Pathway       Nucleus . Nucleus speckle .         Tissue Specificity       Embryo,Heart,Skin,         Function       function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle.,polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population,,similarity. Contains 1 homeobox DNA-binding domain.,subunit:Interacts with RNF 10.,tissue specificity:Embryo and placenta.,         Background       This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a antennapedia-like homeobox-containing genes. The encoded protein may play a	Specificity	MOX-2 Monoclonal Antibody detects endogenous levels of MOX-2 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2 Observed Band 33kD Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo,Heart,Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle. polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population, similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with RNF10.,tissue specificity:Embryo and placenta.,  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo,Heart,Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle.,polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population, similarity.Contains 1 homeobox DNA-binding domain.,subunit:Interacts with RNF10.,tissue specificity:Embryo and placenta.,  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle. polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity:Contains 1 homeobox DNA-binding domain., subunit:Interacts with RNF10., tissue specificity:Embryo and placenta.  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Purification	•
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle. polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity:Contains 1 homeobox DNA-binding domain., subunit:Interacts with RNF10., tissue specificity:Embryo and placenta.,  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Dilution	WB 1:500-2000
Synonyms  MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band  33kD  Cell Pathway  Nucleus . Nucleus speckle .  Tissue Specificity  Embryo, Heart, Skin,  Function  function: Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity: Embryo and placenta.,  Background  This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Concentration	1 mg/ml
Synonyms  MEOX2; GAX; MOX2; Homeobox protein MOX-2; Growth arrest-specific homeobox; Mesenchyme homeobox 2  Observed Band  33kD  Cell Pathway  Nucleus . Nucleus speckle .  Tissue Specificity  Embryo, Heart, Skin,  Function  function: Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity: Embryo and placenta.,  Background  This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Purity	≥90%
homeobox; Mesenchyme homeobox 2  Observed Band 33kD  Cell Pathway Nucleus . Nucleus speckle .  Tissue Specificity Embryo, Heart, Skin,  Function function:Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity:Contains 1 homeobox DNA-binding domain., subunit:Interacts with RNF10., tissue specificity:Embryo and placenta.,  Background This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Storage Stability	-20°C/1 year
Cell Pathway         Nucleus . Nucleus speckle .           Tissue Specificity         Embryo, Heart, Skin,           Function         function: Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity: Embryo and placenta.,           Background         This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Synonyms	·
Tissue Specificity  Embryo, Heart, Skin,  function: Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle., polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population., similarity: Contains 1 homeobox DNA-binding domain., subunit: Interacts with RNF10., tissue specificity: Embryo and placenta.,  Background  This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Observed Band	33kD
function: Role in mesoderm induction and its earliest regional specification, somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle.,polymorphism: The poly-His region of MEOX2 is polymorphic and the number of His varies in the population.,similarity: Contains 1 homeobox DNA-binding domain.,subunit: Interacts with RNF10.,tissue specificity: Embryo and placenta.,  Background  This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Cell Pathway	Nucleus . Nucleus speckle .
somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle.,polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population.,similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with RNF10.,tissue specificity:Embryo and placenta.,  Background  This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like homeobox-containing genes. The encoded protein may play a	Tissue Specificity	Embryo,Heart,Skin,
antennapedia-like homeobox-containing genes. The encoded protein may play a	Function	somitogenesis, and myogenic and sclerotomal differentiation. May have a regulatory role when quiescent vascular smooth muscle cells reenter the cell cycle.,polymorphism:The poly-His region of MEOX2 is polymorphic and the number of His varies in the population.,similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with RNF10.,tissue specificity:Embryo and
	Background	antennapedia-like homeobox-containing genes. The encoded protein may play a

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







mouse protein may be associated with craniofacial and/or skeletal abnormalities, in addition to neurovascular dysfunction observed in Alzheimer's disease. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

