



PAX6 mouse mAb

Catalog No	BYmab-15810
lsotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	PAX6 AN2
Protein Name	PAX6
Immunogen	Synthesized peptide derived from human PAX6
Specificity	This antibody detects endogenous levels of Human PAX6
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Paired box protein Pax-6 (Aniridia type II protein;Oculorhombin)
Observed Band	
Cell Pathway	Nucleus .; [Isoform 1]: Nucleus .; [Isoform 5a]: Nucleus .
Tissue Specificity	[Isoform 1]: Expressed in lymphoblasts. ; [Isoform 5a]: Weakly expressed in lymphoblasts.
Function	developmental stage:Expressed in the developing eye and brain.,disease:Defects in PAX6 are a cause of autosomal dominant keratitis [MIM:148190]. It is an eye disorder characterized by corneal opacification and vascularization, and by foveal hypoplasia.,disease:Defects in PAX6 are a cause of bilateral optic nerve hypoplasia [MIM:165550]; also known as bilateral optic nerve aplasia. Inheritance is autosomal dominant.,disease:Defects in PAX6 are a cause of coloboma of optic nerve [MIM:120430].,disease:Defects in PAX6 are a cause of ectopia pupillae [MIM:129750]. It is a congenital eye malformation in which the pupils are displaced from their normal central position.,disease:Defects in PAX6 are a cause of foveal hypoplasia [MIM:136520]. Foveal hypoplasia can be isolated or associated with presenile cataract. Inheritance is autosomal dominant.,disease:Defects in PAX6 are a cause of Gillespie

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询

Background	This gene encodes a homeobox and paired domain-containing protein that binds DNA and functions as a regulator of transcription. Activity of this protein is key in the development of neural tissues, particularly the eye. This gene is regulated by multiple enhancers located up to hundreds of kilobases distant from this locus. Mutations in this gene or in the enhancer regions can cause ocular disorders such as aniridia and Peter's anomaly. Use of alternate promoters and alternative splicing result in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2015],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd