



EGFL4 Monoclonal Antibody

Catalog No BYmab-15896 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein 8. Function similarity-Contains 12 Kelch repeats, similarity-Contains 2 CUB domains, simi		
Reactivity Human;Rat;Mouse; Applications WB Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse.lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, Function similarity. Contains 12 Kelch repeats, similarity. Contains 2 CUB domains., similarity. Contains 4 laminin EGF-like domains., similarity. Contains 5 EGF-like domains., similarity. Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains., similarity trainst encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains., Similarity trainst encoding different isoforms have been found for this gene.	Catalog No	BYmab-15896
Applications Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, Function similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains, similarity: Contains 0 DEGES in this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, skelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Isotype	IgG
Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, Function similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 4 laminin EGF-like domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two	Reactivity	Human;Rat;Mouse;
Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, domains, similarity: Contains 12 Kelch repeats, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and Psidomains. Defects in this gene are a cause of Carpenter sy	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Pancreas, Plasma, similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and Ps domains. Defects in this gene are a cause of Carpenter syndrome 2. Two	Gene Name	MEGF8
MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, Function similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 4 laminin EGF-like domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Protein Name	Multiple epidermal growth factor-like domains protein 8
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 Iaminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains. The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	lmmunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Specificity	EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains, similarity:Contains 4 laminin EGF-like domains, similarity:Contains 5 EGF-like domains., similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 7 PSI domains., Similarity:Contains 5 EGF-like domains., Similarity:Contains 7 PSI domains., Kelch repeats, and PSI domains. Defects in this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Concentration	1 mg/ml
Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Purity	≥90%
domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Storage Stability	-20°C/1 year
Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 Iaminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Synonyms	domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth
Tissue Specificity Brain,Pancreas,Plasma, similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Observed Band	255kD
Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Cell Pathway	Membrane ; Single-pass type I membrane protein .
domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PS domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Tissue Specificity	Brain,Pancreas,Plasma,
domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.	Function	domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5
	Background	The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using EGFL4 Monoclonal Antibody