



Follistatin Monoclonal Antibody

Catalog No	BYmab-15911
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FST
Protein Name	Follistatin
Immunogen	The antiserum was produced against synthesized peptide derived from human FST. AA range:121-170
Specificity	Follistatin Monoclonal Antibody detects endogenous levels of Follistatin protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FST; Follistatin; FS; Activin-binding protein
Observed Band	65kD
Cell Pathway	Secreted.
Tissue Specificity	Isoform 1 is the predominant isoform in serum but is undetectable in follicular fluid. In the embryo, strong expression is seen in the palatal epithelia, including the medial edge epithelial and midline epithelial seam of the palatal shelves. Less pronounced expression is also seen throughout the palatal shelf and tongue mesenchyme (PubMed:31215115).
Function	function: Binds directly to activin and functions as an activin antagonist. Specific inhibitor of the biosynthesis and secretion of pituitary follicle stimulating hormone (FSH).,similarity: Contains 1 TB (TGF-beta binding) domain.,similarity: Contains 3 follistatin-like domains.,similarity: Contains 3 Kazal-like domains.,subunit: Monomer .,tissue specificity: Isoform 1 is the predominant isoform in serum but is undetectable in follicular fluid.,

Nanjing BYabscience technology Co.,Ltd



Background

Follistatin is a single-chain gonadal protein that specifically inhibits follicle-stimulating hormone release. The single FST gene encodes two isoforms, FST317 and FST344 containing 317 and 344 amino acids respectively, resulting from alternative splicing of the precursor mRNA. In a study in which 37 candidate genes were tested for linkage and association with polycystic ovary syndrome (PCOS) or hyperandrogenemia in 150 families, evidence was found for linkage between PCOS and follistatin. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Follistatin Monoclonal Antibody