



# INSL4 Monoclonal Antibody


Catalog No	BYmab-15942
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	INSL4
Protein Name	Early placenta insulin-like peptide
Immunogen	The antiserum was produced against synthesized peptide derived from human INSL4. AA range:41-90
Specificity	INSL4 Monoclonal Antibody detects endogenous levels of INSL4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	INSL4; Early placenta insulin-like peptide; EPIL; Insulin-like peptide 4; Placentin
Observed Band	19kD
Cell Pathway	Secreted .
Tissue Specificity	Expressed in placenta, uterus and in fetal perichondrium. Expression levels were increased in both early placentas and molar pregnancies and were reduced in choriocarcinoma cells.
Function	developmental stage:Highly expressed in the early placenta. Expression of epil peptides in the villous cytotrophoblast is different from that displayed by the syncytiotrophoblast. In fetal tissues it was identified in the perichondrium of all four limbs, vertebrae, and ribs. It was abundant in interbone ligaments.,function:May play an important role in trophoblast development and in the regulation of bone formation.,similarity:Belongs to the insulin family.,tissue specificity:Expressed in placenta, uterus and in fetal perichondrium. Expression levels were increased in both early placentas and molar pregnancies and were reduced in choriocarcinoma cells.,

Nanjing BYabscience technology Co.,Ltd



<b>Background</b>	INSL4 encodes the insulin-like 4 protein, a member of the insulin superfamily. INSL4 encodes a precursor that undergoes post-translational cleavage to produce 3 polypeptide chains, A-C, that form tertiary structures composed of either all three chains, or just the A and B chains. Expression of INSL4 products occurs within the early placental cytotrophoblast and syncytiotrophoblast. [provided by RefSeq, Jul 2008],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using INSL4 Monoclonal Antibody