



TSG-6 Monoclonal Antibody

Catalog No	BYmab-15975
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TNFAIP6
Protein Name	Tumor necrosis factor-inducible gene 6 protein
Immunogen	The antiserum was produced against synthesized peptide derived from human TSG6. AA range:21-70
Specificity	TSG-6 Monoclonal Antibody detects endogenous levels of TSG-6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TNFAIP6; TSG6; Tumor necrosis factor-inducible gene 6 protein; Hyaluronate-binding protein; TNF-stimulated gene 6 protein; TSG-6; Tumor necrosis factor alpha-induced protein 6; TNF alpha-induced protein 6
Observed Band	31kD
Cell Pathway	Secreted .
Tissue Specificity	Expressed in airway epithelium and submucosal gland (at protein level). Colocalizes with bikunin at the ciliary border. Present in bronchoalveolar lavage fluid (at protein level) (PubMed:16873769). Expressed in mesenchymal stem cells (PubMed:16771708). Found in the synovial fluid of patients with rheumatoid arthritis.
Function	function:Possibly involved in cell-cell and cell-matrix interactions during inflammation and tumorigenesis.,induction:By TNF.,similarity:Contains 1 CUB domain.,similarity:Contains 1 Link domain.,subunit:Interacts with inter-alpha-inhibitor (I-alpha-I). Chondroitin sulfate may be required for the stability of the complex.,tissue specificity:Found in the synovial fluid of patients with rheumatoid arthritis.,

Nanjing BYabscience technology Co.,Ltd



Background

The protein encoded by this gene is a secretory protein that contains a hyaluronan-binding domain, and thus is a member of the hyaluronan-binding protein family. The hyaluronan-binding domain is known to be involved in extracellular matrix stability and cell migration. This protein has been shown to form a stable complex with inter-alpha-inhibitor (I alpha I), and thus enhance the serine protease inhibitory activity of I alpha I, which is important in the protease network associated with inflammation. This gene can be induced by proinflammatory cytokines such as tumor necrosis factor alpha and interleukin-1. Enhanced levels of this protein are found in the synovial fluid of patients with osteoarthritis and rheumatoid arthritis.[provided by RefSeq, Dec 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabs science technology Co.,Ltd