



FGF-9 Monoclonal Antibody

Catalog No	BYmab-16000
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FGF9
Protein Name	Fibroblast growth factor 9
Immunogen	The antiserum was produced against synthesized peptide derived from the C-terminal region of human FGF9. AA range:141-190
Specificity	FGF-9 Monoclonal Antibody detects endogenous levels of FGF-9 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGF9; Fibroblast growth factor 9; FGF-9; Glia-activating factor; GAF; Heparin-binding growth factor 9; HBGF-9
Observed Band	23kD
Cell Pathway	Secreted.
Tissue Specificity	Glial cells.
Function	disease:The continuous overexpression of GAFS may lead to malignant cell growth caused by an autocrine loop.,function:May have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors.,PTM:N-glycosylated.,PTM:Three molecular species were found (30 kDa, 29 kDa and 25 kDa), cleaved at Leu-4, Val-13 and Ser-34 respectively. The smaller ones might be products of proteolytic digestion. Furthermore, there may be a functional signal sequence in the 30 kDa species which is uncleavable in the secretion step.,similarity:Belongs to the heparin-binding growth factors family.,subunit:Monomer.,tissue specificity:Glial

Nanjing BYabscience technology Co.,Ltd



cells.,

Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein was isolated as a secreted factor that exhibits a growth-stimulating effect on cultured glial cells. In nervous system, this protein is produced mainly by neurons and may be important for glial cell development. Expression of the mouse homolog of this gene was found to be dependent on Sonic hedgehog (Shh) signaling. Mice lacking the homolog gene displayed a male-to-female sex reversal phenotype, which suggested a role in testicular embryogenesis. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using FGF-9 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658