



RhoGEF p115 Monoclonal Antibody

Catalog No	BYmab-16247
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ARHGEF1
Protein Name	Rho guanine nucleotide exchange factor 1
Immunogen	The antiserum was produced against synthesized peptide derived from human ARHGEF1. AA range:162-211
Specificity	RhoGEF p115 Monoclonal Antibody detects endogenous levels of RhoGEF p115 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ARHGEF1; Rho guanine nucleotide exchange factor 1; 115 kDa guanine nucleotide exchange factor; p115-RhoGEF; p115RhoGEF; Sub1.5
Observed Band	105kD
Cell Pathway	Cytoplasm . Membrane . Translocated to the membrane by activated GNA13 or LPA stimulation.
Tissue Specificity	Ubiquitously expressed.
Function	domain:The DH domain is involved in interaction with CCPG1.,domain:The RGS domain, also known as rgRGS domain, is necessary but not sufficient for GAP activity.,function:Seems to play a role in the regulation of RhoA GTPase by guanine nucleotide-binding alpha-12 (GNA12) and alpha-13 (GNA13) subunits. Acts as GTPase-activating protein (GAP) for GNA12 and GNA13, and as guanine nucleotide exchange factor (GEF) for RhoA GTPase. Activated G alpha 13/GNA13 stimulates the RhoGEF activity through interaction with the RGS-like domain. This GEF activity is inhibited by binding to activated GNA12.,PTM:Phosphorylated by PKCA.,sequence caution:Contaminating sequence. Sequence of unknown origin in the N-terminal part.,similarity:Contains

Nanjing BYabscience technology Co.,Ltd



1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 RGS (RGS-like) domain.,subcellular location:Translocated to the membrane by activ

Background

Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form complex with G proteins and stimulate Rho-dependent signals. Multiple alternatively spliced transcript variants have been found for this gene, but the full-length nature of some variants has not been defined. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using RhoGEF p115 Monoclonal Antibody