

# CaV $\alpha$ 2 $\delta$ 2 Monoclonal Antibody

<b>Catalog No</b>	BYmab-16323
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	CACNA2D2
<b>Protein Name</b>	Voltage-dependent calcium channel subunit alpha-2/delta-2 (Voltage-gated calcium channel subunit alpha-2/delta-2) [Cleaved into: Voltage-dependent calcium channel subunit alpha-2-2; Voltage-dependent
<b>Immunogen</b>	Synthetic Peptide of CaV $\alpha$ 2 $\delta$ 2 AA range: 540-620
<b>Specificity</b>	The antibody detects endogenous CaV $\alpha$ 2 $\delta$ 2 protein
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Voltage-dependent calcium channel subunit alpha-2/delta-2 (Voltage-gated calcium channel subunit alpha-2/delta-2) [Cleaved into: Voltage-dependent calcium channel subunit alpha-2-2; Voltage-dependent calcium channel subunit delta-2]
<b>Observed Band</b>	100-120kD
<b>Cell Pathway</b>	Membrane ; Single-pass type I membrane protein . Colocalizes with CACNA1A in lipid raft fractions. .
<b>Tissue Specificity</b>	Predominantly present in cerebellar cortex. Present in various lung tumor cell lines, while it is absent in normal lung (at protein level). Highly expressed in heart, lung, testis, pancreas and skeletal muscle. Also expressed in kidney, liver, placenta and brain.
<b>Function</b>	domain:The MIDAS-like motif in the VWFA domain binds divalent metal cations and is required to promote trafficking of the alpha-1 (CACNA1) subunit to the plasma membrane by an integrin-like switch.,function:The alpha-2/delta subunit of voltage-dependent calcium channels regulates calcium current density and

**Nanjing BYabs science technology Co.,Ltd**



activation/inactivation kinetics of the calcium channel. Acts as a regulatory subunit for P/Q-type calcium channel (CACNA1A), N-type (CACNA1B), L-type (CACNA1C OR CACNA1D) and possibly T-type (CACNA1G). Overexpression induces apoptosis.,miscellaneous: Binds gabapentin, an antiepileptic drug.,PTM: May be proteolytically processed into subunits alpha-2-2 and delta-2 that are disulfide-linked. It is however unclear whether such cleavage really takes place in vivo and has a functional role.,PTM: N-glycosylated.,similarity: Belongs to the calcium channel subunit alpha-2/delta family.,similarit

## Background

calcium voltage-gated channel auxiliary subunit alpha2delta 2(CACNA2D2) Homo sapiens Calcium channels mediate the entry of calcium ions into the cell upon membrane polarization. This gene encodes the alpha-2/delta subunit of the voltage-dependent calcium channel complex. The complex consists of the main channel-forming subunit alpha-1, and auxiliary subunits alpha-2/delta, beta, and gamma. The auxiliary subunits function in the assembly and membrane localization of the complex, and modulate calcium currents and channel activation/inactivation kinetics. The subunit encoded by this gene undergoes post-translational cleavage to yield the extracellular alpha2 peptide and a membrane-anchored delta polypeptide. This subunit is a receptor for the antiepileptic drug, gabapentin. Mutations in this gene are associated with early infantile epileptic encephalopathy. Single nucleotide polymorphisms in this gene are correlated with increased sensitivity to


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using CaV  $\alpha$  2  $\delta$  2 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd