



CLC-KA Monoclonal Antibody

| Catalog No | BYmab-16403 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | CLCNKA |
| Protein Name | Chloride channel protein CIC-Ka |
| Immunogen | The antiserum was produced against synthesized peptide derived from human CLCNKA. AA range:581-630 |
| Specificity | CLC-KA Monoclonal Antibody detects endogenous levels of CLC-KA protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | CLCNKA; Chloride channel protein ClC-Ka; Chloride channel Ka; ClC-K1 |
| Observed Band | 75kD |
| Cell Pathway | Membrane; Multi-pass membrane protein. |
| Tissue Specificity | Expressed predominantly in the kidney. All nephron segments expressing BSND also express CLCNK proteins. |
| Function | function:Voltage-gated chloride channel. Chloride channels have several functions including the regulation of cell volume; membrane potential stabilization, signal transduction and transepithelial transport. May be important in urinary concentrating mechanisms.,similarity:Belongs to the chloride channel (TC 2.A.49) family.,similarity:Contains 2 CBS domains.,subunit:Interacts with BSND. Forms heteromers with BSND in the thin ascending limb of Henle.,tissue specificity:Expressed predominantly in the kidney. All nephron segments expressing BSND also express CLCNK proreins., |
| Background | This gene is a member of the CLC family of voltage-gated chloride channels. The encoded protein is predicted to have 12 transmembrane domains, and requires a |
| | |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| | beta subunit called barttin to form a functional channel. It is thought to function in salt reabsorption in the kidney and potassium recycling in the inner ear. The gene is highly similar to CLCNKB, which is located 10 kb downstream from this gene. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Western Blot analysis of various cells using CLC-KA Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658