



GlyRβ Monoclonal Antibody

| Catalog No | BYmab-16428 |
|--------------------|---|
| lsotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | GLRB |
| Protein Name | Glycine receptor subunit beta |
| Immunogen | The antiserum was produced against synthesized peptide derived from human GLRB. AA range:211-260 |
| Specificity | GlyR $\beta~$ Monoclonal Antibody detects endogenous levels of GlyR $\beta~$ protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | GLRB; Glycine receptor subunit beta; Glycine receptor 58 kDa subunit |
| Observed Band | 56kD |
| Cell Pathway | Cell junction, synapse, postsynaptic cell membrane ; Multi-pass membrane protein . Cell junction, synapse . Cell projection, dendrite . Cell membrane ; Multi-pass membrane protein . Cytoplasm . Retained in the cytoplasm upon heterologous expression by itself. Coexpression with GPHN promotes expression at the cell membrane (PubMed:12684523). Coexpression with GLRA1, GLRA2 or GLRA3 promotes expression at the cell membrane . |
| Tissue Specificity | Brain,Hippocampus, |
| Function | disease:Defects in GLRB are a cause of startle disease (STHE) [MIM:149400]; also known as hereditary hyperekplexia or congenital stiff-person syndrome. STHE is a genetically heterogeneous neurologic disorder characterized by muscular rigidity of central nervous system origin, particularly in the neonatal period, and by an exaggerated startle response to unexpected acoustic or tactile |
| | stimuli. Inheritance can be autosomal dominant or recessive., function: The glycine receptor is a neurotransmitter-gated ion channel. Binding of glycine to its receptor |

| 博研生物 BYabscience | 国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询 |
|----------------------------|---|
| | increases the chloride conductance and thus produces hyperpolarization (inhibition of neuronal firing).,similarity:Belongs to the ligand-gated ionic channel (TC 1.A.9) family.,subunit:Pentamer composed of alpha and beta subunits. Interacts with GPHN., |
| Background | This gene encodes the beta subunit of the glycine receptor, which is a pentamer composed of alpha and beta subunits. The receptor functions as a neurotransmitter-gated ion channel, which produces hyperpolarization via increased chloride conductance due to the binding of glycine to the receptor. Mutations in this gene cause startle disease, also known as hereditary hyperekplexia or congenital stiff-person syndrome, a disease characterized by muscular rigidity. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009], |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Western Blot analysis of various cells using GlyR β Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd