



HCN2 Monoclonal Antibody

Catalog No	BYmab-16429
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HCN2
Protein Name	Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2
Immunogen	The antiserum was produced against synthesized peptide derived from human HCN2. AA range:491-540
Specificity	HCN2 Monoclonal Antibody detects endogenous levels of HCN2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HCN2; BCNG2; Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2; Brain cyclic nucleotide-gated channel 2; BCNG-2
Observed Band	100kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Highly expressed throughout the brain. Detected at low levels in heart.
Function	domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position.,function:Hyperpolarization-activated ion channel exhibiting weak selectivity for potassium over sodium ions. Contributes to the native pacemaker currents in heart (If) and in neurons (Ih). Produces a large instantaneous current. Activated by cAMP. Modulated by intracellular chloride ions and pH; acidic pH shifts the activation to more negative voltages.,miscellaneous:Inhibited by extracellular cesium ions.,similarity:Belongs to the potassium channel HCN family.,similarity:Contains 1 cyclic nucleotide-binding domain.,subunit:The potassium channel is probably composed of a homo- or heterotetrameric complex

Nanjing BYabscience technology Co.,Ltd



of pore-forming subunits. Heteromultimer with HCN1. Interacts with KCNE2.,tissue specificity:Highly expressed throughout the brain. Detected at

Background

Hyperpolarization-activated cation channels of the HCN gene family, such as HCN2, contribute to spontaneous rhythmic activity in both heart and brain.[supplied by OMIM, Jul 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using HCN2 Monoclonal Antibody