



NMDAε4 Monoclonal Antibody

Catalog No	BYmab-16489
Isotype	IgG
Reactivity	Human;Mouse;Rat;Monkey
Applications	WB
Gene Name	GRIN2D
Protein Name	Glutamate [NMDA] receptor subunit epsilon-4
Immunogen	The antiserum was produced against synthesized peptide derived from human GRIN2D. AA range:671-720
Specificity	NMDA ϵ 4 Monoclonal Antibody detects endogenous levels of NMDA ϵ 4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GRIN2D; GluN2D; NMDAR2D; Glutamate [NMDA] receptor subunit epsilon-4; EB11; N-methyl D-aspartate receptor subtype 2D; NMDAR2D; NR2D
Observed Band	170kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein.
Tissue Specificity	Brain,Fetal brain,
Function	function:NMDA receptor subtype of glutamate-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Mediated by glycine.,similarity:Belongs to the glutamate-gated ion channel (TC 1.A.10) family.,subunit:Interacts with PDZ domains of INADL and DLG4 (By similarity). Forms heteromeric channel of a zeta subunit (GRIN1), a epsilon subunit (GRIN2A, GRIN2B, GRIN2C or GRIN2D) and a third subunit (GRIN3A or GRIN3B).,
Background	N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA channel has been shown to be involved in long-term

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



potentiation, an activity-dependent increase in the efficiency of synaptic
transmission thought to underlie certain kinds of memory and learning. NMDA
receptor channels are heteromers composed of the key receptor subunit
NMDAR1 (GRIN1) and 1 or more of the 4 NMDAR2 subunits: NMDAR2A
(GRIN2A), NMDAR2B (GRIN2B), NMDAR2C (GRIN2C), and NMDAR2D
(GRIN2D). [provided by RefSeq, Mar 2010],
(GININZD). [provided by Nersey, Mar 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NMDA ϵ 4 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd