



ASIC1 Monoclonal Antibody

Catalog No	BYmab-16534
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ASIC1 ACCN2 BNAC2
Protein Name	Acid-sensing ion channel 1 (ASIC1) (Amiloride-sensitive cation channel 2, neuronal) (Brain sodium channel 2) (BNAc2)
Immunogen	Synthetic peptide from human protein at AA range: 220-280
Specificity	The antibody detects endogenous ASIC1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Acid-sensing ion channel 1 (ASIC1;Amiloride-sensitive cation channel 2, neuronal;Brain sodium channel 2;BNAc2)
Observed Band	55kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Localizes in synaptosomes at dendritic synapses of neurons. Colocalizes with DLG4 (By similarity). .
Tissue Specificity	Expressed in most or all neurons.
Function	alternative products:The splice variant from ASIC1a described in mouse and rat, which gives rise to an isoform with different N-termini (Asic1b), does not seem to exist in human,function:Cation channel with high affinity for sodium, which is gated by extracellular protons and inhibited by the diuretic amiloride. Also permeable for Ca(2+), Li(+) and K(+). Generates a biphasic current with a fast inactivating and a slow sustained phase. Mediates glutamate-independent Ca(2+) entry into neurons upon acidosis. This Ca(2+) overloading is toxic for cortical neurons and may be in part responsible for ischemic brain injury. Heteromeric channel assembly seems to modulate channel properties. Functions as a postsynaptic proton receptor that influences intracellular Ca(2+) concentration and calmodulin-dependent protein kinase II phosphorylation and thereby the

Nanjing BYabscience technology Co.,Ltd



density of dendritic spines. Modulates a

Background

This gene encodes a member of the acid-sensing ion channel (ASIC) family of proteins, which are part of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. Members of the ASIC family are sensitive to amiloride and function in neurotransmission. The encoded proteins function in learning, pain transduction, touch sensation, and development of memory and fear. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ASIC1 Monoclonal Antibody