



MCM4 Monoclonal Antibody

Catalog No	BYmab-16750
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	MCM4
Protein Name	DNA replication licensing factor MCM4
Immunogen	The antiserum was produced against synthesized peptide derived from human MCM4. AA range:20-69
Specificity	MCM4 Monoclonal Antibody detects endogenous levels of MCM4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MCM4; CDC21; DNA replication licensing factor MCM4; CDC21 homolog; P1-CDC21
Observed Band	96kD
Cell Pathway	Nucleus . Chromosome . Associated with chromatin before the formation of nuclei and detaches from it as DNA replication progresses
Tissue Specificity	Cervix,Epithelium,Testis,
Function	function:Involved in the control of DNA replication.,similarity:Belongs to the MCM family.,similarity:Contains 1 MCM domain.,
Background	The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 6 and 7 proteins possesses DNA helicase activity, and may act as a DNA

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

精准的 WB 检测服务 (3)。

24H 在线服务,欢迎咨询



unwir the D is ma	ding enzyme. The phosphorylation of this protein by CDC2 kinase reduces IA helicase activity and chromatin binding of the MCM complex. This gene ped to a region on the chromosome 8 head-to-head next to the C/DNA-PK, a DNA-activated protein kinase involved in the repair of DNA
PRKI doub	C/DNA-PK, a DNA-activated protein kinase involved in the repair of DNA e-strand breaks. Alternatively spliced transcri
matters needing Avoid attention	repeated freezing and thawing!
Usage suggestions This more	roduct can be used in immunological reaction related experiments. For nformation, please consult technical personnel.
	Products Images
3T3 K562	3T3 Western Blot analysis of various cells using MCM4 Monoclonal Antibody 117 85 48 34 26 19 (kD)

Nanjing BYabscience technology Co.,Ltd