



POT1 Monoclonal Antibody

BYmab-16769
IgG
Human;Monkey
WB
POT1
Protection of telomeres protein 1
The antiserum was produced against synthesized peptide derived from human POT1. AA range:247-296
POT1 Monoclonal Antibody detects endogenous levels of POT1 protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
POT1; Protection of telomeres protein 1; hPot1; POT1-like telomere end-binding protein
70kD
Nucleus . Chromosome, telomere . Colocalizes with telomeric DNA.
Ubiquitous.
alternative products:A number of isoforms are produced, function:Component of the telomerase ribonucleoprotein (RNP) complex that is essential for the replication of chromosome termini. Is a component of the double-stranded telomeric DNA-binding TRF1 complex which is involved in the regulation of telomere length by cis-inhibition of telomerase. Also acts as a single-stranded telomeric DNA-binding protein and thus may act as a downstream effector of the TRF1 complex and may transduce information about telomere maintenance and/or length to the telomere terminus. Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. Shelterin associates with arrays of double-stranded TTAGGG repeats added by

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



telomerase and protects chromosome ends; without its p	rotective activity.
telomeres are no longer hidden from the DNA damage s	urveillance and

Background

protection of telomeres 1(POT1) Homo sapiens This gene is a member of the telombin family and encodes a nuclear protein involved in telomere maintenance. Specifically, this protein functions as a member of a multi-protein complex that binds to the TTAGGG repeats of telomeres, regulating telomere length and protecting chromosome ends from illegitimate recombination, catastrophic chromosome instability, and abnormal chromosome segregation. Increased transcriptional expression of this gene is associated with stomach carcinogenesis and its progression. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using POT1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd