



Cadherin-16 Monoclonal Antibody

Catalog No	BYmab-16927
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CDH16
Protein Name	Cadherin-16
Immunogen	The antiserum was produced against synthesized peptide derived from human CDH16. AA range:780-829
Specificity	Cadherin-16 Monoclonal Antibody detects endogenous levels of Cadherin-16 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CDH16; Cadherin-16; Kidney-specific cadherin; Ksp-cadherin
Observed Band	90kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein .
Tissue Specificity	Kidney specific.
Function	function:Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells cadherins may thus contribute to the sorting of heterogeneous cell types.,similarity:Contains 6 cadherin domains.,tissue specificity:Kidney specific.,
Background	This gene is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-associated glycoproteins. Mapped to a previously identified cluster of cadherin genes on chromosome 16q22.1, the gene localizes with superfamily members CDH1, CDH3, CDH5, CDH8 and CDH11. The protein consists of an extracellular domain containing 6 cadherin domains, a transmembrane region and a truncated cytoplasmic domain but lacks the
	Naniing Byahscience technology Co. Ltd

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商


精准的 WB 检测服务

24H 在线服务, 欢迎咨询



	prosequence and tripeptide HAV adhesion recognition sequence typical of most classical cadherins. Expression is exclusively in kidney, where the protein functions as the principal mediator of homotypic cellular recognition, playing a role in the morphogenic direction of tissue development. Alternatively spliced transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Mar 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd