



# COL19A1 Monoclonal Antibody

<b>Catalog No</b>	BYmab-16976
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	COL19A1
<b>Protein Name</b>	Collagen alpha-1(XIX) chain
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human Collagen XIX alpha1. AA range:421-470
<b>Specificity</b>	COL19A1 Monoclonal Antibody detects endogenous levels of COL19A1 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	COL19A1; Collagen alpha-1(XIX) chain; Collagen alpha-1(Y) chain
<b>Observed Band</b>	120kD
<b>Cell Pathway</b>	Secreted, extracellular space, extracellular matrix .
<b>Tissue Specificity</b>	Localized to vascular, neuronal, mesenchymal, and some epithelial basement membrane zones in umbilical cord.
<b>Function</b>	domain:The numerous interruptions in the triple helix may make this molecule either elastic or flexible.,function:May act as a cross-bridge between fibrils and other extracellular matrix molecules. Involved in skeletal myogenesis in the developing esophagus. May play a role in organization of the pericellular matrix or the sphinteric smooth muscle.,PTM:Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.,similarity:Belongs to the fibril-associated collagens with interrupted helices (FACIT) family.,similarity:Contains 1 TSP N-terminal (TSPN) domain.,subunit:Oligomer; disulfide-linked.,tissue specificity:Localized to vascular, neuronal, mesenchymal, and some epithelial basement membrane zones in umbilical cord.,

**Nanjing BYabscience technology Co.,Ltd**

**Background**

This gene encodes the alpha chain of type XIX collagen, a member of the FACIT collagen family (fibril-associated collagens with interrupted helices). Although the function of this collagen is not known, other members of this collagen family are found in association with fibril-forming collagens such as type I and II, and serve to maintain the integrity of the extracellular matrix. The transcript produced from this gene has an unusually large 3' UTR which has not been completely sequenced. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images