



# Laminin $\alpha$ -1 Monoclonal Antibody

|                    |  |
|--------------------|--|
| Catalog No         | BYmab-17037  |
| Isotype            | IgG  |
| Reactivity         | Human;Rat;Mouse;   |
| Applications       | WB   |
| Gene Name          | LAMA1  |
| Protein Name       | Laminin subunit alpha-1  |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human LAMA1. AA range:2501-2550  |
| Specificity        | Laminin $\alpha$ -1 Monoclonal Antibody detects endogenous levels of Laminin $\alpha$ -1 protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Monoclonal, Mouse,IgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-2000  |
| Concentration      | 1 mg/ml  |
| Purity             | $\geq 90\%$  |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | LAMA1; LAMA; Laminin subunit alpha-1; Laminin A chain; Laminin-1 subunit alpha; Laminin-3 subunit alpha; S-laminin subunit alpha; S-LAM alpha  |
| Observed Band      |  |
| Cell Pathway       | Secreted, extracellular space, extracellular matrix, basement membrane. Major component.   |
| Tissue Specificity | Liver,Placenta,Skin,   |
| Function           | domain:Domains VI, IV and G are globular.,domain:The alpha-helical domains I and II are thought to interact with other laminin chains to form a coiled coil structure.,function:Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components.,similarity:Contains 1 laminin N-terminal domain.,similarity:Contains 17 laminin EGF-like domains.,similarity:Contains 2 laminin IV type A domains.,similarity:Contains 5 laminin G-like domains.,subcellular location:Major component.,subunit:Laminin is a complex glycoprotein, consisting of three different polypeptide chains (alpha, beta, gamma), which are bound to each other |

**Nanjing BYabscience technology Co.,Ltd**



by disulfide bonds into a cross-shaped molecule comprising one long and three short arms with globules at each end. Alp

#### Background

This gene encodes one of the alpha 1 subunits of laminin. The laminins are a family of extracellular matrix glycoproteins that have a heterotrimeric structure consisting of an alpha, beta and gamma chain. These proteins make up a major component of the basement membrane and have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Mutations in this gene may be associated with Poretti-Boltshauser syndrome. [provided by RefSeq, Sep 2014],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Western Blot analysis of various cells using Laminin  $\alpha$ -1 Monoclonal Antibody