




## GAB2 (Phospho Tyr614) mouse mAb

| Catalog No         | BYmab-17191  |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human, Mouse,Rat   |
| Applications       | WB   |
| Gene Name          | GAB2 KIAA0571  |
| Protein Name       | GRB2-associated-binding protein 2 (GRB2-associated binder 2) (Growth factor receptor bound protein 2-associated protein 2) (pp100)   |
| Immunogen          | Synthesized peptide derived from human GAB2 (Phospho Tyr614)   |
| Specificity        | This antibody detects endogenous levels of GAB2 (Phospho Tyr614) Mouse mAb at Human, Mouse,Rat   |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| Source             | Mouse,Monoclonal   |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | GRB2-associated-binding protein 2 (GRB2-associated binder 2) (Growth factor receptor bound protein 2-associated protein 2) (pp100)   |
| Observed Band      | 100kD  |
| Cell Pathway       | Cytoplasm . Cell membrane .  |
| Tissue Specificity |  |
| Function           | PTM:Dephosphorylated by PTPN11.,PTM:Phosphorylated on tyrosine residue(s) by the thrombopoietin receptor (TPOR), stem cell factor receptor (SCFR), and T-cell and B-cell antigen receptors, gp130, IL-2R and IL-3R.,similarity:Belongs to the GAB family.,similarity:Contains 1 PH domain.,subunit:Interacts with GRB2, PI-3 kinase and with other SH2-containing proteins., |
| Background         | GRB2 associated binding protein 2(GAB2) Homo sapiens This gene is a member of the GRB2-associated binding protein (GAB) gene family. These proteins contain pleckstrin homology (PH) domain, and bind SHP2 tyrosine phosphatase and GRB2 adapter protein. They act as adapters for transmitting various signals in response to stimuli through cytokine and growth factor    |
|                    |  |

Nanjing BYabscience technology Co.,Ltd








|                           | receptors, and T- and B-cell antigen receptors. The protein encoded by this gene is the principal activator of phosphatidylinositol-3 kinase in response to activation of the high affinity IgE receptor. Two alternatively spliced transcripts encoding different isoforms have been described for this gene. [provided by RefSeq, Nov 2009], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing!   |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |

## **Products Images**



Western Blot analysis of various cells using GAB2 (Phospho Tyr614) mouse mAb

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658