



K1199 mouse mAb

Catalog No	BYmab-17232
Isotype	IgG
Reactivity	Human, Mouse
Applications	WB
Gene Name	KIAA1199
Protein Name	Protein KIAA1199
Immunogen	Synthesized peptide derived from human C-terminal K1199
Specificity	This antibody detects endogenous levels of K1199 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Mouse, Monoclonal
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Protein KIAA1199
Observed Band	
Cell Pathway	Nucleus. Cytoplasm. Endoplasmic reticulum. Cell membrane. Membrane, clathrin-coated pit. Secreted. Retained in the endoplasmic reticulum (ER) in a HSPA5/BIP-dependent manner. Colocalized with clathrin heavy chain/CLTC in clathrin-coated vesicles. Strongly detected in the cytoplasm of breast carcinoma cells, whereas poorly detected in adjacent normal epithelial cells, stromal cells, or benign breast tissues. Localized in the nucleus and cytoplasm of colon adenocarcinomas.
Tissue Specificity	Expressed in dermal and in synovial fibroblasts. Strongly expressed in gastric cancers compared with the paired normal tissues. Strongly expressed in both ductal carcinoma and invasive breast cancer cells compared with benign epithelial cells (at protein level). Strongly expressed in brain, placenta, prostate, breast, lung and testis. Expressed in fibroblasts, epithelial cells and cancer cells. In ear, it is specifically expressed in inner ear. Expressed in cochlea and vestibule tissues. Strongly expressed in gastric cancers compared with the paired normal tissues. Strongly expressed in colon adenocarcinomas compared with normal colonic mucosas. Strongly expressed in breast cancer as compared to normal breast

Nanjing BYabscience technology Co.,Ltd



tissue.

Function

Mediates depolymerization of hyaluronic acid (HA) via the cell membrane-associated clathrin-coated pit endocytic pathway. Binds to hyaluronic acid. Hydrolyzes high molecular weight hyaluronic acid to produce an intermediate-sized product, a process that may occur through rapid vesicle endocytosis and recycling without intracytoplasmic accumulation or digestion in lysosomes. Involved in hyaluronan catabolism in the dermis of the skin and arthritic synovium. Positively regulates epithelial-mesenchymal transition (EMT), and hence tumor cell growth, invasion and cancer dissemination. In collaboration with HSPA5/BIP, promotes cancer cell migration in a calcium and PKC-dependent manner. May be involved in hearing.

Background**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658