



VATL mouse mAb

Catalog No	BYmab-17242
Isotype	IgG
Reactivity	Human, Mouse,Rat
Applications	WB
Gene Name	ATP6V0C ATP6C ATP6L ATPL
Protein Name	V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit)
Immunogen	Synthesized peptide derived from human N-ternal VATL
Specificity	This antibody detects endogenous levels of VATL at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Mouse,Monoclonal
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit)
	V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid
Synonyms	V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit)
Synonyms Observed Band	 V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit) 17kD Cytoplasmic vesicle, clathrin-coated vesicle membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ;
Synonyms Observed Band Cell Pathway	 V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit) 17kD Cytoplasmic vesicle, clathrin-coated vesicle membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Multi-pass membrane protein . Proton-conducting pore forming of the V0 complex of vacuolar(H+)-ATPase (V-ATPase), a multisubunit enzyme composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons . V-ATPase is responsible for acidifying and maintaining the pH of intracellular
Synonyms Observed Band Cell Pathway Tissue Specificity	 V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit) 17kD Cytoplasmic vesicle, clathrin-coated vesicle membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Multi-pass membrane protein . Proton-conducting pore forming of the V0 complex of vacuolar(H+)-ATPase (V-ATPase), a multisubunit enzyme composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons . V-ATPase is responsible for acidifying and maintaining the pH of intracellular compartments and in some cell types. is targeted to the plasma membrane, where

Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd