



## WWP1 mouse mAb

Catalog No	BYmab-17276
Isotype	IgG
Reactivity	Human, Mouse
Applications	WB
Gene Name	WWP1
Protein Name	NEDD4-like E3 ubiquitin-protein ligase WWP1 (EC 6.3.2) (Atrophin-1-interacting protein 5) (AIP5) (TGIF-interacting ubiquitin ligase 1) (Tiul1) (WW domain-containing protein 1)
Immunogen	Synthesized peptide derived from human N-ternal WWP1
Specificity	This antibody detects endogenous levels of WWP1 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Mouse,Monoclonal
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NEDD4-like E3 ubiquitin-protein ligase WWP1 (EC 6.3.2) (Atrophin-1-interacting protein 5) (AIP5) (TGIF-interacting ubiquitin ligase 1) (Tiul1) (WW domain-containing protein 1)
Observed Band	
Cell Pathway	Cytoplasm . Cell membrane ; Peripheral membrane protein . Nucleus .
Tissue Specificity	Detected in heart, placenta, pancreas, kidney, liver, skeletal muscle, bone marrow, fetal brain, and at much lower levels in adult brain and lung. Isoform 1 and isoform 5 predominate in all tissues tested, except in testis and bone marrow, where isoform 5 is expressed at much higher levels than isoform 1.
Function	E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Ubiquitinates ERBB4 isoforms JM-A CYT-1 and JM-B CYT-1, KLF2, KLF5 and TP63 and promotes their proteasomal degradation. Ubiquitinates RNF11 without targeting it for degradation. Ubiquitinates and promotes degradation of TGFBR1; the ubiquitination is
	Naniing BYabscience technology CoLtd

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务,欢迎咨询

	enhanced by SMAD7. Ubiquitinates SMAD6 and SMAD7. Ubiquitinates and promotes degradation of SMAD2 in response to TGF-beta signaling, which requires interaction with TGIF.
Background	
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
	Products Images

Nanjing BYabscience technology Co.,Ltd