



## OGDHL mouse mAb

Catalog No	BYmab-17283
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	OGDHL KIAA1290
Protein Name	2-oxoglutarate dehydrogenase-like, mitochondrial (EC 1.2.4) (2-oxoglutarate dehydrogenase complex component E1-like) (OGDC-E1-like) (Alpha-ketoglutarate dehydrogenase-like)
Immunogen	Synthesized peptide derived from human C-ternal OGDHL
Specificity	This antibody detects endogenous levels of OGDHL at Human
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Mouse, Monoclonal
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	2-oxoglutarate dehydrogenase-like, mitochondrial (EC 1.2.4) (2-oxoglutarate dehydrogenase complex component E1-like) (OGDC-E1-like) (Alpha-ketoglutarate dehydrogenase-like)
Observed Band	
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	Expressed in the brain and the liver.
Function	2-oxoglutarate dehydrogenase (E1-like) component of the 2-oxoglutarate dehydrogenase multienzyme complex (OGDHC) which mediates the decarboxylation of alpha-ketoglutarate in the tricarboxylic acid cycle. The OGDHC complex catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO(2) while reducing NAD(+) to NADH (By similarity). The OGDHC complex is mainly active in the mitochondrion (By similarity). Involved in the inhibition of cell proliferation and in apoptosis.

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background
------------

matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**