



ODO2 mouse mAb

Catalog No	BYmab-17290
Isotype	lgG
Reactivity	Human, Mouse,Rat
Applications	WB
Gene Name	DLST DLTS
Protein Name	Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolip
Immunogen	Synthesized peptide derived from human C-ternal ODO2
Specificity	This antibody detects endogenous levels of ODO2 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Mouse,Monoclonal
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex) (E2K)
Observed Band	
Cell Pathway	Mitochondrion matrix . Nucleus . Mainly localizes in the mitochondrion. A small fraction localizes to the nucleus, where the 2-oxoglutarate dehydrogenase complex is required for histone succinylation
Tissue Specificity	
Function	Dihydrolipoamide succinyltransferase (E2) component of the 2-oxoglutarate dehydrogenase complex. The 2-oxoglutarate dehydrogenase complex catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO(2). The 2-oxoglutarate dehydrogenase complex is mainly active in the mitochondrion . A fraction of the 2-oxoglutarate dehydrogenase complex also localizes in the nucleus and is required for lysine succinylation of histones: associates with
	Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Products Images		
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
matters needing attention	Avoid repeated freezing and thawing!	
Background		
	KAT2A on chromatin and provides succinyl-CoA to histone succinyltransferase KAT2A .	

Nanjing BYabscience technology Co.,Ltd