






Toll-Like Receptor 4 mouse mAb

| Catalog No | BYmab-17807 |
|--|---|
| Isotype | IgG |
| Reactivity | Human |
| Applications | WB |
| Gene Name | TLR4 |
| Protein Name | |
| Immunogen | TLR4; Toll-like receptor 4; hToll; CD antigen CD284 |
| Specificity | |
| Formulation | |
| Source | |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| | |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% -20°C/1 year |
| | |
| Storage Stability | -20°C/1 year |
| Storage Stability Synonyms | -20°C/1 year TLR4; Toll-like receptor 4; hToll; CD antigen CD284 |
| Storage Stability Synonyms Observed Band | -20°C/1 year TLR4; Toll-like receptor 4; hToll; CD antigen CD284 |
| Storage Stability Synonyms Observed Band Cell Pathway | -20°C/1 year TLR4; Toll-like receptor 4; hToll; CD antigen CD284 |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | -20°C/1 year TLR4; Toll-like receptor 4; hToll; CD antigen CD284 |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | -20°C/1 year TLR4; Toll-like receptor 4; hToll; CD antigen CD284 Calculated MW: 96 kDa; Observed MW: 90-120 kDa Cooperates with LY96 and CD14 to mediate the innate immune response to |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | -20°C/1 year TLR4; Toll-like receptor 4; hToll; CD antigen CD284 Calculated MW: 96 kDa; Observed MW: 90-120 kDa Cooperates with LY96 and CD14 to mediate the innate immune response to bacterial lipopolysaccharide (LPS). |

Nanjing BYabscience technology Co.,Ltd







Products Images

Nanjing BYabscience technology Co.,Ltd