




SET mouse mAb

Catalog No BYmab-17842 Isotype IgG Reactivity Human,Mouse,Rat Applications WB Gene Name SET Protein Name Formulation Specificity Formulation Source Formulation Burification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding, Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. Wasse suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.		
Reactivity Human,Mouse,Rat Applications WB Gene Name SET Protein Name Immunogen A synthesized peptide derived from human SET Specificity Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Catalog No	BYmab-17842
Applications Gene Name SET Protein Name Immunogen A synthesized peptide derived from human SET Specificity Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Isotype	IgG
Gene Name Protein Name Immunogen A synthesized peptide derived from human SET Specificity Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histore binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. Matters needing attention This product can be used in immunological reaction related experiments. For	Reactivity	Human,Mouse,Rat
Protein Name Immunogen A synthesized peptide derived from human SET Specificity Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. Matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Applications	WB
Immunogen A synthesized peptide derived from human SET Specificity Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Gene Name	SET
Specificity Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Protein Name	
Formulation Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; 12PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding, Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Immunogen	A synthesized peptide derived from human SET
Source Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Avoid repeated freezing and thawing! Usage suggestions This product can be used in immunological reaction related experiments. For	Specificity	
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Formulation	
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Source	
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention This product can be used in immunological reaction related experiments. For	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Usage suggestions This product can be used in immunological reaction related experiments. For	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention This product can be used in immunological reaction related experiments. For	Concentration	1 mg/ml
Synonyms 2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Avoid repeated freezing and thawing! Usage suggestions This product can be used in immunological reaction related experiments. For	Purity	≥90%
Observed Band Calculated MW: 33 kDa; Observed MW: 39 kDa Cell Pathway Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention This product can be used in immunological reaction related experiments. For	Storage Stability	-20°C/1 year
Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Avoid repeated freezing and thawing! This product can be used in immunological reaction related experiments. For	Synonyms	2PP2A; IGAAD; TAF-I; I2PP2A; IPP2A2; PHAPII; TAF-IBETA
Tissue Specificity Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Avoid repeated freezing and thawing! Usage suggestions This product can be used in immunological reaction related experiments. For	Observed Band	Calculated MW: 33 kDa; Observed MW: 39 kDa
Function Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. Matters needing attention Avoid repeated freezing and thawing! Usage suggestions This product can be used in immunological reaction related experiments. For	Cell Pathway	
Background Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. Avoid repeated freezing and thawing! Usage suggestions This product can be used in immunological reaction related experiments. For	Tissue Specificity	
and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. matters needing attention Avoid repeated freezing and thawing! This product can be used in immunological reaction related experiments. For	Function	
Usage suggestionsThis product can be used in immunological reaction related experiments. For	Background	Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone binding. Isoform 2 anti-apoptotic activity is mediated by inhibition of
Usage suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.		the GZMA-activated DNase, NME1.
	matters needing attention	

Nanjing BYabscience technology Co.,Ltd







Products Images

Nanjing BYabscience technology Co.,Ltd