



## LC3A/B mouse mAb

Catalog No	BYmab-17896
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	MAP1LC3A/MAP1LC3B
Protein Name	
Immunogen	Synthetic peptide.
Specificity	
Formulation	
Source	
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3; MAP1LC3A; LC3B; ATG8F; MAP1LC3B-a; MAP1A/1BLC3; MAP1LC3B
Observed Band	Calculated MW: 14 kDa; Observed MW: 14,16 kDa
Cell Pathway	
Tissue Specificity	
Function	
Background	Macroautophagy is the major inducible pathway for the general turnover of

Nanjing BYabscience technology Co.,Ltd

cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome

(or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
matters needing attention	Avoid repeated freezing and thawing!
	microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd