



CENPE mouse mAb

Catalog No	BYmab-17900
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	CENPE
Protein Name	Centromere-associated protein E (Centromere protein E) (CENP-E) (Kinesin-related protein CENPE)
Immunogen	Synthesized peptide derived from human CENPE
Specificity	This antibody detects endogenous levels of CENPE at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	297kD
Cell Pathway	Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle . Chromosome, centromere . Associates with kinetochores during congression (as early as prometaphase), relocates to the spindle midzone at anaphase, and is quantitatively discarded at the end of the cell division (By similarity). Recruited to the kinetochore in a SEPT7, CENPQ and TRAPPC12-dependent manner (PubMed:18460473, PubMed:25918224, PubMed:25395579). Recruited to the pericentromeric/centromeric regions of the chromosome in a CTCF-dependent manner (PubMed:26321640).
Tissue Specificity	
Function	Microtubule plus-end-directed kinetochore motor which plays an important role in chromosome congression, microtubule-kinetochore conjugation and spindle assembly checkpoint activation. Drives chromosome congression (alignment of chromosomes at the spindle equator resulting in the formation of the metaphase plate) by mediating the lateral sliding of polar chromosomes along spindle

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	microtubules towards the spindle equator and by aiding the establishment and maintenance of connections between kinetochores and spindle microtubules. The transport of pole-proximal chromosomes towards the spindle equator is favored by microtubule tracks that are detyrosinated. Acts as a processive bi-directional tracker of dynamic microtubule tips; after chromosomes have congressed, continues to play an active role at kinetochores, enhancing their links with dynamic microtubule ends. Suppresses chromosome c
Background	
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658