



SLC40A1 mouse mAb

| Catalog No | BYmab-17948 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | SLC40A1 FPN1 IREG1 SLC11A3 MSTP079 |
| Protein Name | Solute carrier family 40 member 1 (Ferroportin-1) (Iron-regulated transporter 1) |
| Immunogen | Synthesized peptide derived from human SLC40A1. AA range 1-100 |
| Specificity | This antibody detects endogenous levels of SLC40A1 at Human, Mouse,Rat |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 63kD |
| Cell Pathway | Cell membrane ; Multi-pass membrane protein . Localized to the basolateral membrane of polarized epithelial cells. |
| Tissue Specificity | Detected in erythrocytes (at protein level). Expressed in placenta, intestine, muscle and spleen. |
| Function | Major iron transporter that plays a key role in balancing cellular and systemic iron levels. Transports iron from intestinal, splenic, and hepatic cells into the blood to provide iron to other tissues (By similarity). Controls therefore dietary iron uptake, iron recycling by macrophages, and release of iron stores in hepatocytes (By similarity). When iron is in excess, hepcidin/HAMP levels increase resulting in a degradation of ferroportin/SLC40A1 limiting the iron efflux to plasma. |
| Background | |

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd