



RHOC mouse mAb

Catalog No	BYmab-18065
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	RHOC ARH9 ARHC
Protein Name	Rho-related GTP-binding protein RhoC (Rho cDNA clone 9) (h9)
Immunogen	Synthesized peptide derived from human RHOC
Specificity	This antibody detects endogenous levels of RHOC at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year 21kD
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 21kD Cell membrane ; Lipid-anchor ; Cytoplasmic side . Cleavage furrow . Translocates
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 21kD Cell membrane ; Lipid-anchor ; Cytoplasmic side . Cleavage furrow . Translocates
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 21kD Cell membrane ; Lipid-anchor ; Cytoplasmic side . Cleavage furrow . Translocates to the equatorial region before furrow formation in a ECT2-dependent manner. Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Serves as a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Regulates apical junction formation in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 21kD Cell membrane ; Lipid-anchor ; Cytoplasmic side . Cleavage furrow . Translocates to the equatorial region before furrow formation in a ECT2-dependent manner. Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Serves as a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Regulates apical junction formation in

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd